arXiv:1401.3992v1 [math.RA] 16 Jan 2014

Classification of five dimensional nilpotent Jordan algebras

Ahmed Hegazi, Hani Abdelwahab
Mathematics department, Faculty of Science, Mansoura University, Egypt
hegazi@mans.edu.eg
hanii_i@yahoo.com

January 17, 2014

Abstract

We use the method which has been described in [1] to give a classification of 5-dimensional nilpotent Jordan algebras over an algebraically closed fields of characteristic $\neq 2$. We show that there are 44 classes of non-associative nilpotent Jordan algebras over an algebraically closed fields of characteristic $\neq 2$, up to isomorphism each class has only one algebra except 6 classes contain infinite number of non-isomorphic algebras.

Keywords and phrases: *Nilpotent Jordan algebras, second cohomology, centeral extension, automorphism group, orbits, isomorphism.*
2010 Mathematics Subject Classification: *17C10, 17C55, 17-08*

1 Introduction

The classification of small-dimensional Jordan algebras is a classical problem. The authors described a computional method to construct nilpotent Jordan algebras over any field and used this method to classify nilpotent Jordan algebras of dimensions $\leqslant 3$ over any field and four dimensional nilpotent algebras over an algebraically closed fields of characteristic $\neq 2$ and real field $\mathbb{R}$ (see [1]). The original aim of the research presented in this paper was to extend the results to 5-dimensional nilpotent Jordan algebras over an algebraically closed fields of characteristic $\neq 2$. Nilpotent associative Jordan algebras of dimension five were classified by Mazzola ([2] - nilpotent commutative associative algebras of dimension $\leqslant 5$, over algebraically closed fields of characteristic not 2,3), and recently, Poonen ([3] - nilpotent commutative associative algebras of dimension $\leqslant 5$, over

algebraically closed fields). To complete the classification, we construct all 5-dimensional nilpotent non-associative Jordan algebras over an algebraically closed fields of characteristic $\neq 2$. Our methodology, which is the same as in [1] with some restrictions to avoid reconstructing commutative associative algebras is explained in Section 2.

Here is an outline of the paper. In Section 2 we describe the cohomological method that we use to classify nilpotent Jordan algebras, which also appeared in [1]. In Section 3 we present the main results of [1], that is the classification of the nilpotent Jordan algebras of dimension $\leqslant 4$ which are needed in Section 4. Then in Section 4 the main work is performed to classify the 5-dimensional nilpotent non-associative Jordan algebras, some results relied on computer calculations (specifically, computing a Gröbner bases for ideals in a polynomial rings) and were inspired by Gröbner basis computations in the computational algebra system Magma [4]. Finally in section 5 we summarize our main results of *classification of five dimensional nilpotent Jordan algebras*.

Remark 1.1. Throughout the paper $\mathbb{K}$ be an algebraically closed fields of characteristic $\neq 2$. Also we use some notational conventions, if J be Jordan algebra with basis elements $a_1, \ldots, a_n$. Then by δ_{a_i,a_j} we denote the symmetric bilinear map $J \times J \to \mathbb{K}$ with $\delta_{a_i,a_j}(a_k, a_l) = 1$ if $i = k$ and $j = l$, or $i = l$ and$j = k$, and it takes the value 0 otherwise.

2 A summary of the method

The main idea that we use here is to obtain the nilpotent Jordan algebras of dimension n as central extensions of Jordan algebras of smaller dimension. The central extensions are defined using the second cohomology space, and the isomorphism classes of the central extensions correspond to the orbits of the automorphism group on the set of the subspaces of this cohomology space.

Let J be a Jordan algebra, for a vector space V, let $Z^2_{Jor.}(J, V)$ denote the set of symmetric bilinear maps $\theta : J \times J \to V$ with the property that

$$\theta(a, d\circ(b\circ c)) + \theta(b, d\circ(a\circ c)) + \theta(c, d\circ(a\circ b)) = \theta(a\circ b, c\circ d) + \theta(b\circ c, a\circ d) + \theta(a\circ c, b\circ d)$$

for all $a, b, c, d \in J$. The set $Z^2_{Jor.}(J, V)$ is viewed as a vector space over $\Bbbk$ and the elements of $Z^2_{Jor.}(J, V)$ are said to be *Jordan cocycles.* A linear map $f : J \longrightarrow V$ is called a 1-*cochain* from J to V. The set of all 1-*cochains* from J to V is denoted by $C^1(J, V)$. Then the bilinear map $\delta f : J \times J \longrightarrow V$defined by $(\delta f)(x, y) := f(x \circ y)$ is a *Jordan cocycle,* called a *coboundary* from J to V. The set of all *coboundaries* is denoted by $\delta C^1(J, V)$, and the *second cohomology space* $H^2_{Jor.}(J, V)$ is defined as the quotient $Z^2_{Jor.}(J, V)/\delta C^1(J, V)$.

Let J be a Jordan algebra and V a vector space over a field $\Bbbk$, we denote the center of J by $Z(J)$. For $\theta \in Z^2_{Jor.}(J, V)$, define a Jordan algebra J_θ as follows. The underlying space of J_θ is $J \oplus V$. The product of two elements $a + v,\ b + w \in J_\theta$, is defined as $(a + v) \circ\ (b + w) = a \circ_J b + \theta(a, b)$ where $a \circ_J b$ denotes the product in J. Then J_θ is

a Jordan algebra and V is an ideal of J_θ such that $V \leqslant Z(J_\theta)$. In addition, $J \cong J_\theta/V$, and hence J_θ is a central extension of J. Further, if $\theta_1,\ \theta_2 \in Z^2_{Jor.}(J,V)$ such that $\theta_1 - \theta_2 \in \delta C^1(J,V)$ then $J_{\theta_1} \cong J_{\theta_2}$, and so the isomorphism type of J_θ only depends on the element $\theta + \delta C^1(J,V)$ of $H^2_{Jor.}(J,V)$. Conversely let M be a Jordan algebra such that $Z(M) \neq 0$, and set $V = Z(M)$ and $J = M/Z(M)$. Let $\pi\colon M \to J$ be the projection map. Choose an injective linear map $f\colon J \to M$ such that $\pi(f(a)) = a$ for all $a \in J$. Define $\theta\colon J \times J \to V$ by $\theta(a,b) = f(a \circ b) - f(a) \circ f(b)$. Then θ is a cocycle such that $M \cong J_\theta$. Though θ depends on the choice of f, the coset $\theta + \delta C^1(J,V)$ is independent of f. Hence the central extension M of J determines a well-defined element of $H^2_{Jor.}(J,V)$.

Let V be m-dimensional vector space then $Z^2_{Jor.}(J,V) = Z^2_{Jor.}(J,\Bbbk^m) = Z^2_{Jor.}(J,\Bbbk)^m$ and $H^2_{Jor.}(J,V) = H^2_{Jor.}(J,\Bbbk^m) = H^2_{Jor.}(J,\Bbbk)^m$. Let us now fix a basis $\{e_1,\ldots,e_m\}$ of V. A Jordan cocycle $\theta \in Z^2_{Jor.}(J,V)$ can be written as

$$\theta(a,b) = \sum_{i=1}^{m} \theta_i(a,b)e_i,$$

where $\theta_i \in Z^2_{Jor.}(J,\Bbbk)$. Furthermore, θ is a coboundary if and only if all θ_i are. For $\theta \in Z^2(J,V)$, let $\theta^\perp$ denote the *radical* of θ; that is, the set of elements $a \in J$ such that $\theta(a,b) = 0$ for all $b \in J$. Then

$$\theta^\perp = \theta_1^\perp \cap \theta_2^\perp \cap .... \cap \theta_m^\perp.$$

For each $\theta \in Z^2_{Jor.}(J,V)$ and $\phi \in Aut(J)$, the automorphism group of J. Define $\phi\theta(a,b) = \theta(\phi(a),\phi(b))$ for any $a,b \in J$. So $Aut(J)$ acts on $Z^2_{Jor.}(J,V)$, and $\phi\theta \in \delta C^1(J,V)$ if and only if $\theta \in \delta C^1(J,V)$. Then $Z^2_{Jor.}(J,V)$ and $\delta C^1(J,V)$ are invariant under the action of $Aut(J)$. So $Aut(J)$ acts on $H^2_{Jor.}(J,V)$ hence the quotient $H^2_{Jor.}(J,V)$ can be viewed as an $Aut(J)$-module.

Lemma 2.1 (see [1]). *Let $\theta = (\theta_1,\theta_2,.....,\theta_m)$ and $\eta = (\eta_1,\eta_2,.....,\eta_m) \in H^2_{Jor.}(J,V)$ and $\theta^\perp \cap Z(J) = \eta^\perp \cap Z(J) = 0$. Then J_θ and J_η are isomorphic if and only if there exist $\phi \in Aut(J)$ such that $\phi\theta_i$ span the same subspace of $H^2_{Jor.}(J,\Bbbk)$ as the η_i .*

A Jordan algebra J is said to be that J has a central component if $J = J_1 \oplus \Bbbk$ where J_1 is an ideal of J and $\Bbbk$ is viewed as a 1-dimensional Jordan algebra, in this case J can be viewed as 1-dimensional centeral extension of J_1 corresponding to $\theta = 0$ " Trivial 1-dimensional centeral extension".

Lemma 2.2 (see [1]). *Let $\theta(a,b) = \sum_{i=1}^{m} \theta_i(a,b)e_i \in H^2_{Jor.}(J,V)$ and $\theta^\perp \cap Z(J) = 0$. Then J_θ has a centeral component if and only if $\theta_1,\theta_2,.....,\theta_m$ are linearly dependent .*

Note that $J = J_1 \oplus \Bbbk$ is associative if and only if J_1 is associative, So if we need to construct all nilpotent non-associative Jordan algebras of dimension n with centeral components we will only consider the trivial 1-dimensional centeral extension of all nilpotent non-associative Jordan algebras of dimension $n-1$.

Let $G_r(H^2(J,\Bbbk))$ be the Grassmanian of subspaces of dimension r in $H^2(J,\Bbbk)$. There is anatural action of $Aut(J)$ on $G_r(H^2(J,\Bbbk))$ defined by :

$$\phi < \theta_1, \theta_2, ..., \theta_r >=< \phi\theta_1, \phi\theta_2, ..., \phi\theta_r >$$

for $V =< \theta_1, \theta_2, ..., \theta_r > \in G_r(H^2(J,\Bbbk))$ and $\phi \in Aut(J)$. Define

$$U_r(J) = \{V =< \theta_1, \theta_2, ..., \theta_r > \in G_r(H^2(J,\Bbbk)) : \theta_i^{\perp} \cap Z(J) = 0; i = 1, 2, ..., r\}.$$

Theorem 2.3 (see [1]). *Let $U_r(J)/Aut(J)$ be the set of $Aut(J)$-orbits of $U_r(J)$. Then there exists a canonical one-to-one correspondense from $U_r(J)/Aut(J)$ onto the set of isomorphism classes of Jordan algebras without centeral components which are central extensions of J by V and have r-dimensional center where $r = \dim V$.*

Now to avoid construction of associative nilpotent Jordan algebras, define a subspace $H^2_{Ass.}(J,V)$ of $H^2_{Jor.}(J,V)$ as follows :

$$H^2_{Ass.}(J,V) = \{\theta \in H^2_{Jor.}(J,V) : \theta(a \circ b, c) = \theta(a, b \circ c) \text{ for all } a, b, c \in J\}.$$

Then J_θ is associative if and only if J is associative and $\theta \in H^2_{Ass.}(J,V)$, conversly J_θ is non-associative if and only if J is non-associative or J is associative and $\theta \notin H^2_{Ass.}(J,V)$. We will call θ a non-associative Jordan cocycle if $\theta \in H^2_{Jor.}(J,V)$ and $\theta \notin H^2_{Ass.}(J,V)$.

All nilpotent Jordan algebras of dimensions $\leqslant 3$ are associative. Hence the classification of 5-dimensional nilpotent non-associative Jordan algebras requires that we determine the trivial and non-trivial centeral extension of nilpotent non-associative algebras of dimension 4, non-trivial centeral extension of nilpotent associative algebras of dimension at most 4 by non-associative Jordan cocycles. The determination of these algebras is achieved in Section 4.

3 Nilpotent Jordan algebras of dimension $\leqslant 4$

Here and in next sections the basis elements of the algebras will be denoted by the letters $a, b, \ldots$. The multiplication of an algebra is specified by giving only the nonzero products among the basis elements.We denote the j-th algebra of dimension i by $J_{i,j}$.

There are the following nilpotent Jordan algebras of dimensions $\leqslant 4$ (cf. e.g., [1]):

Algebra	Multiplication taple	$Z(J)$	Comments
$J_{1,1}$	$-\ -\ -\ -\ -\ -$	a	associative
$J_{2,1}$	$-\ -\ -\ -\ -\ -$	a, b	associative
$J_{2,2}$	$a^2 = b$	b	associative
$J_{3,1}$	$-\ -\ -\ -\ -\ -$	a, b, c	associative
$J_{3,2}$	$a^2 = b$	b, c	associative
$J_{3,3}$	$a \circ b = c$	c	associative
$J_{3,4}$	$a^2 = b, a \circ b = c$	c	associative
$J_{4,1}$	$-\ -\ -\ -\ -\ -$	a, b, c, d	associative
$J_{4,2}$	$a^2 = b$	b, c, d	associative
$J_{4,3}$	$a \circ b = c$	c, d	associative
$J_{4,4}$	$a^2 = b, a \circ b = c$	c, d	associative
$J_{4,5}$	$a \circ b = d, c^2 = d$	d	associative
$J_{4,6}$	$a^2 = b\ ,\ b \circ c = d$	d	non-associative
$J_{4,7}$	$a^2 = b\ ,\ a \circ b = d\ ,\ c^2 = d$	d	associative
$J_{4,8}$	$a \circ b = c, a \circ c = d$	d	non-associative
$J_{4,9}$	$a \circ b = c, a \circ c = d, b^2 = d$	d	non-associative
$J_{4,10}$	$a \circ b = c, a \circ c = d, b \circ c = d$	d	non-associative
$J_{4,11}$	$a^2 = b\ ,\ a \circ b = c\ ,\ a \circ c = d\ ,\ b^2 = d$	d	associative
$J_{4,12}$	$a^2 = c\ ,\ a \circ b = d$	c, d	associative
$J_{4,13}$	$a^2 = c\ ,\ b^2 = d$	c, d	associative

Remark 3.1. We replaced under isomorphism some certain algebras in [1] by $J_{4,3}$, $J_{4,5}$, $J_{4,8}$, $J_{4,9}$, $J_{4,10}$ and $J_{4,13}$ introducing as they were in the previous table. To be somewhat more concrete, let J_1, J_2, J_3, J_4, J_5 and J_6 are four dimensional nilpotent Jordan algebras defined as follow

- $J_1 : a'^2 = c', b'^2 = c'$.
- $J_2 : a'^2 = c', b'^2 = c', a' \circ c' = d'$.
- $J_3 : a'^2 = c', b'^2 = -c', a' \circ c' = d', b' \circ c' = d'$.
- $J_4 : a'^2 = c', b'^2 = -c', a' \circ c' = d', b' \circ c' = d', a' \circ b' = d'$.
- $J_5 : a'^2 = d', b'^2 = d', c'^2 = d'$.
- $J_6 : a'^2 = c', b'^2 = c', a' \circ b' = d'$.

Define $\varphi_1, \varphi_2, \varphi_3, \varphi_4$ and φ_5 as follow :

- $\varphi_1(a') = a + \frac{b}{2}, \varphi_1(b') = \sigma\left(a - \frac{b}{2}\right), \varphi_1(c') = c, \varphi_1(d') = d$ where $\sigma^2 = -1$, then $J_1 \cong J_{4,3}$ and $J_5 \cong J_{4,5}$.
- $\varphi_2(a') = \frac{a+b}{2}, \varphi_2(b') = \frac{a-b}{2}, \varphi_2(c') = \frac{1}{2}c, \varphi_2(d') = \frac{1}{2}d$, then $J_2 \cong J_{4,10}$.
- $\varphi_3(a') = a + \frac{b}{2}, \varphi_3(b') = a - \frac{b}{2}, \varphi_3(c') = c, \varphi_3(d') = d$, then $J_3 \cong J_{4,8}$.
- $\varphi_4(a') = a - b - \frac{c}{2}, \varphi_4(b') = a + b - \frac{c}{2}, \varphi_4(c') = -2c, \varphi_4(d') = -2d$, then $J_4 \cong J_{4,9}$.
- $\varphi_5(a') = a + b, \varphi_5(b') = a - b, \varphi_5(c') = c + d, \varphi_5(d') = c - d$, then $J_6 \cong J_{4,13}$.

Now we describe $H^2_{Ass.}(J, \Bbbk)$ only for nilpotent associative Jordan algebras and $H^2_{Jor.}(J, \Bbbk)$ for both associative and non-associative nilpotent Jordan algebras of dimension $\leqslant 4$ to determine algebras and their cocycles whose centeral extensions are nilpotent non-associative Jordan algebras. We get the following :

Algebra	$H^2_{Ass.}(J, \Bbbk)$	$H^2_{Jor.}(J, \Bbbk)$
$J_{1,1}$	$< \delta_{a,a} >$	$H^2_{Ass.}(J_{1,1}, \Bbbk)$
$J_{2,1}$	$< \delta_{a,a}, \delta_{a,b}, \delta_{b,b} >$	$H^2_{Ass.}(J_{2,1}, \Bbbk)$
$J_{2,2}$	$< \delta_{a,b} >$	$H^2_{Ass.}(J_{2,2}, \Bbbk)$
$J_{3,1}$	$< \delta_{a,a}, \delta_{b,b}, \delta_{c,c}, \delta_{a,b}, \delta_{a,c}, \delta_{b,c} >$	$H^2_{Ass.}(J_{3,1}, \Bbbk)$
$J_{3,2}$	$< \delta_{a,b}, \delta_{a,c}, \delta_{c,c} >$	$H^2_{Ass.}(J_{3,2}, \Bbbk) \oplus < \delta_{b,c} >$
$J_{3,3}$	$< \delta_{a,a}, \delta_{b,b} >$	$H^2_{Ass.}(J_{3,2}, \Bbbk) \oplus < \delta_{a,c}, \delta_{b,c} >$
$J_{3,4}$	$< \delta_{a,c} + \delta_{b,b} >$	$H^2_{Ass.}(J_{3,4}, \Bbbk)$
$J_{4,1}$	$< \delta_{a,a}, \delta_{b,b}, \delta_{c,c}, \delta_{d,d}, \delta_{a,b}, \delta_{a,c}, \delta_{a,d}, \delta_{b,c}, \delta_{b,d}, \delta_{c,d} >$	$H^2_{Ass.}(J_{4,1}, \Bbbk)$
$J_{4,2}$	$< \delta_{c,c}, \delta_{d,d}, \delta_{a,b}, \delta_{a,c}, \delta_{a,d}, \delta_{c,d} >$	$H^2_{Ass.}(J_{4,2}, \Bbbk) \oplus < \delta_{b,c}, \delta_{b,d} >$
$J_{4,3}$	$< \delta_{a,a}, \delta_{b,b}, \delta_{d,d}, \delta_{a,d}, \delta_{b,d} >$	$H^2_{Ass.}(J_{4,3}, \Bbbk) \oplus < \delta_{a,c}, \delta_{b,c}, \delta_{c,d} >$
$J_{4,4}$	$< \delta_{a,c} + \delta_{b,b}, \delta_{a,d}, \delta_{d,d} >$	$H^2_{Ass.}(J_{4,4}, \Bbbk) \oplus < \delta_{b,d} >$
$J_{4,5}$	$< \delta_{a,a}, \delta_{b,b}, \delta_{c,c}, \delta_{a,b}, \delta_{a,c}, \delta_{b,c} > / < \delta_{c,c} + \delta_{a,b} >$	$H^2_{Ass.}(J_{4,4}, \Bbbk) \oplus < \delta_{a,d}, \delta_{b,d}, \delta_{c,d} >$
$J_{4,6}$	$- - - - - -$	$< \delta_{a,b}, \delta_{a,c}, \delta_{c,c}, \delta_{b,d} >$
$J_{4,7}$	$< \delta_{a,c}, \delta_{a,b}, \delta_{c,c} > / < \delta_{a,b} + \delta_{c,c} >$	$H^2_{Ass.}(J_{4,7}, \Bbbk) \oplus < \delta_{b,b} + \delta_{a,d} >$
$J_{4,8}$	$- - - - - -$	$< \delta_{a,a}, \delta_{b,b}, \delta_{b,c} >$
$J_{4,9}$	$- - - - - -$	$< \delta_{a,a}, \delta_{b,b}, \delta_{a,c}, \delta_{b,c} > / < \delta_{b,b} + \delta_{a,c} >$
$J_{4,10}$	$- - - - - -$	$< \delta_{a,a}, \delta_{b,b}, \delta_{a,c}, \delta_{b,c} > / < \delta_{a,c} + \delta_{b,c} >$
$J_{4,11}$	$< \delta_{a,d} + \delta_{b,b} >$	$H^2_{Ass.}(J_{4,11}, \Bbbk)$
$J_{4,12}$	$< \delta_{a,c}, \delta_{b,b}, \delta_{a,d} + \delta_{b,c} >$	$< \delta_{a,c}, \delta_{b,b}, \delta_{a,d}, \delta_{b,c}, \delta_{b,d} >$
$J_{4,13}$	$< \delta_{a,b}, \delta_{a,c}, \delta_{b,d} >$	$H^2_{Ass.}(J_{4,13}, \Bbbk) \oplus < \delta_{b,c}, \delta_{a,d} >$

From the previous table we have the following lemma:

Lemma 3.2. *Every nilpotent non-associative Jordan algebra of dimension five is one of the following :*

1. *1-dimensional centeral extension of nilpotent non-associative Jordan algebras $J_{4,6}, J_{4,8}, J_{4,9}$ and $J_{4,10}$.*
2. *1-dimensional centeral extension of $J_{4,2}, J_{4,3}, J_{4,4}, J_{4,5,}J_{4,7}, J_{4,12}$ and $J_{4,13}$ by non-associative Jordan cocycles.*
3. *2-dimensional centeral extension of $J_{3,2}$ and $J_{3,3}$ by non-associative Jordan cocycles.*

4 Non-associative nilpotent Jordan algebras

In this section we compute centeral extensions for algebras stated in lemma 3.2. We have a number of subsections; in each subsection the central extensions of one particular Jordan algebra are considered excpet at subsection 4.2 we consider Jordan algebras $J_{4,8,}J_{4,9}$ and $J_{4,10}$.

4.1 1-dimensional centeral extension of $J_{4,6}$

The Jordan algebra $J_{4,6}$ is non-associative, firstly we consider 1-dimensional trivial centeral extension of $J_{4,6}$ corresponding to $\theta = 0$. We get algebra :

- $J_{5,1} = J_{4,6} \oplus J_{1,1}$.

Next we consider non-trivial 1-dimensional centeral extension of $J_{4,6}$. Let $\theta = \alpha_1\delta_{a,b} + \alpha_2\delta_{a,c} + \alpha_3\delta_{b,d} + \alpha_4\delta_{c,c} \in H^2_{Jor.}(J_{4,6}, \Bbbk)$, the center of $J_{4,6}$, $Z(J_{4,6})$, is spanned by d. Furthermore the automorphism group, $Aut(J_{4,6})$, consists of :

$$\phi = \begin{bmatrix} a_{11} & 0 & 0 & 0 \\ a_{21} & a_{11}^2 & 0 & 0 \\ a_{31} & 0 & a_{33} & 0 \\ a_{41} & 2a_{21}a_{31} & a_{43} & a_{11}^2a_{33} \end{bmatrix}, a_{11}a_{33} \neq 0.$$

The automorphism group acts as follows :
$\alpha_1 \longrightarrow a_{11}^2\left(\alpha_1a_{11} + \alpha_3a_{41}\right) + 2\alpha_3a_{21}^2a_{31}$,
$\alpha_2 \longrightarrow a_{33}\left(\alpha_2a_{11} + \alpha_4a_{31}\right) + \alpha_3a_{21}a_{43}$,
$\alpha_3 \longrightarrow \alpha_3a_{11}^4a_{33}$,
$\alpha_4 \longrightarrow \alpha_4a_{33}^2$.

We need $\alpha_3 \neq 0$ in order to have that d do not lie in the radical of θ. Choose $a_{31} = 0, a_{41} = -\frac{\alpha_1a_{11}}{\alpha_3}, a_{21} = a_{11}a_{33}$ and $a_{43} = -\frac{1}{\alpha_3}$ we obtain that $[\alpha_1, \alpha_2, \alpha_3, \alpha_4] \longrightarrow [0, 0, \alpha_3a_{11}^4a_{33}, \alpha_4a_{33}^2]$. If $\alpha_4 = 0$, choose $a_{11} = 1$ and $a_{33} = \frac{1}{\alpha_3}$ we get a cocycle $\theta_1 = \delta_{b,d}$,

on the other hand if $\alpha_4 \neq 0$, choose $a_{11} = \sqrt[4]{\frac{\sqrt{\alpha_4}}{\alpha_3}}$ and $a_{33} = \frac{1}{\sqrt{\alpha_4}}$ we get a cocycle $\theta_2 = \delta_{b,d} + \delta_{c,c}$. It easy to see that θ_1 and θ_2 are not in the same $Aut(J_{4,6})$-orbit, so we get two algebras :

- $J_{5,2} : a^2 = b, b \circ c = d, b \circ d = e.$
- $J_{5,3} : a^2 = b, b \circ c = d, b \circ d = e, c^2 = e.$

4.2 1-dimensional centeral extension of $J_{4,8,}J_{4,9}$ and $J_{4,10}$.

The Jordan algebras $J_{4,8,}J_{4,9}$ and $J_{4,10}$ are non-associative and have no non-trivial 1-dimensional centeral extensions since the radical of $\theta's$ always intersect with the center of these algebras. So we only consider 1-dimensional trivial centeral extensions of $J_{4,8,}J_{4,9}$ and $J_{4,10}$. We get algebras :

- $J_{5,4} = J_{4,8} \oplus J_{1,1}$.
- $J_{5,5} = J_{4,9} \oplus J_{1,1}$.
- $J_{5,6} = J_{4,10} \oplus J_{1,1}$.

4.3 1-dimensional centeral extension of $J_{4,2}$

Let $\theta = \alpha_1\delta_{c,c} + \alpha_2\delta_{d,d} + \alpha_3\delta_{a,b} + \alpha_4\delta_{a,c} + \alpha_5\delta_{a,d} + \alpha_6\delta_{c,d} + \alpha_7\delta_{b,c} + \alpha_8\delta_{b,d}$ such that $(\alpha_7, \alpha_8) \neq (0, 0)$. The center of

$J_{4,2}$, $Z(J_{4,2})$, is spanned by b, c and d. Furthermore the automorphism group, $Aut(J_{4,2})$, consists of :

$$\phi = \begin{bmatrix} a_{11} & 0 & 0 & 0 \\ a_{21} & a_{11}^2 & a_{23} & a_{24} \\ a_{31} & 0 & a_{33} & a_{34} \\ a_{41} & 0 & a_{43} & a_{44} \end{bmatrix}, a_{11}\left(a_{33}a_{44} - a_{34}a_{43}\right) \neq 0.$$

The automorphism group acts as follows :

$$\begin{aligned}
\alpha_1 &\longrightarrow a_{33}\left(\alpha_1 a_{33} + \alpha_7 a_{23} + \alpha_6 a_{43}\right) + a_{43}\left(\alpha_2 a_{43} + \alpha_6 a_{33} + \alpha_8 a_{23}\right) + a_{23}\left(\alpha_7 a_{33} + \alpha_8 a_{43}\right) \\
\alpha_2 &\longrightarrow a_{34}\left(\alpha_1 a_{34} + \alpha_7 a_{24} + \alpha_6 a_{44}\right) + a_{44}\left(\alpha_2 a_{44} + \alpha_6 a_{34} + \alpha_8 a_{24}\right) + a_{24}\left(\alpha_7 a_{34} + \alpha_8 a_{44}\right) \\
\alpha_3 &\longrightarrow a_{11}^2\left(\alpha_3 a_{11} + \alpha_7 a_{31} + \alpha_8 a_{41}\right) \\
\alpha_4 &\longrightarrow a_{33}\left(\alpha_1 a_{31} + \alpha_4 a_{11} + \alpha_7 a_{21} + \alpha_6 a_{41}\right) + a_{43}\left(\alpha_2 a_{41} + \alpha_5 a_{11} + \alpha_6 a_{31} + \alpha_8 a_{21}\right) \\
&\quad + a_{23}\left(\alpha_3 a_{11} + \alpha_7 a_{31} + \alpha_8 a_{41}\right) \\
\alpha_5 &\longrightarrow a_{34}\left(\alpha_1 a_{31} + \alpha_4 a_{11} + \alpha_7 a_{21} + \alpha_6 a_{41}\right) + a_{44}\left(\alpha_2 a_{41} + \alpha_5 a_{11} + \alpha_6 a_{31} + \alpha_8 a_{21}\right) \\
&\quad + a_{24}\left(\alpha_3 a_{11} + \alpha_7 a_{31} + \alpha_8 a_{41}\right) \\
\alpha_6 &\longrightarrow a_{34}\left(\alpha_1 a_{33} + \alpha_7 a_{23} + \alpha_6 a_{43}\right) + a_{44}\left(\alpha_2 a_{43} + \alpha_6 a_{33} + \alpha_8 a_{23}\right) + a_{24}\left(\alpha_7 a_{33} + \alpha_8 a_{43}\right)
\end{aligned}$$

$\alpha_7 \longrightarrow a_{11}^2 (\alpha_7 a_{33} + \alpha_8 a_{43})$
$\alpha_8 \longrightarrow a_{11}^2 (\alpha_7 a_{34} + \alpha_8 a_{44})$

No generality is lost by assuming that $(\alpha_7, \alpha_8) = (1, 0)$. To fix $(\alpha_7, \alpha_8) \longrightarrow (1, 0)$ choose $a_{11} = a_{33} = 1$ and $a_{34} = 0$ then :

$\alpha_1 \longrightarrow \alpha_1 + 2a_{23} + a_{43} (\alpha_2 a_{43} + 2\alpha_6)$
$\alpha_2 \longrightarrow \alpha_2 a_{44}^2$
$\alpha_3 \longrightarrow \alpha_3 + a_{31}$
$\alpha_4 \longrightarrow (\alpha_1 a_{31} + \alpha_4 + a_{21} + \alpha_6 a_{41}) + a_{43} (\alpha_2 a_{41} + \alpha_5 + \alpha_6 a_{31}) + a_{23} (\alpha_3 + a_{31})$
$\alpha_5 \longrightarrow a_{44} (\alpha_2 a_{41} + \alpha_5 + \alpha_6 a_{31}) + a_{24} (\alpha_3 + a_{31})$
$\alpha_6 \longrightarrow a_{44} (\alpha_2 a_{43} + \alpha_6) + a_{24}$
$\alpha_7 \longrightarrow 1$
$\alpha_8 \longrightarrow 0$

Choose $a_{43} = 0, a_{23} = -\frac{1}{2}\alpha_1, a_{31} = -\alpha_3$ and $a_{24} = -\alpha_6 a_{44}$ we obtain that :
$\alpha_1 \longrightarrow 0, \alpha_3 \longrightarrow 0, \alpha_6 \longrightarrow 0, \alpha_7 \longrightarrow 1, \alpha_8 \longrightarrow 0,$
$\alpha_2 \longrightarrow \alpha_2 a_{44}^2,$
$\alpha_4 \longrightarrow -\alpha_1 \alpha_3 + \alpha_4 + a_{21} + \alpha_6 a_{41},$
$\alpha_5 \longrightarrow a_{44} (\alpha_2 a_{41} + \alpha_5 - \alpha_3 \alpha_6).$

Thus no generality is lost by assuming that $[\alpha_1, \alpha_3, \alpha_6, \alpha_7, \alpha_8] = [0, 0, 0, 1, 0]$.

To fix $[\alpha_1, \alpha_3, \alpha_6, \alpha_7, \alpha_8] \longrightarrow [0, 0, 0, 1, 0]$ take $a_{11} = a_{33} = 1$ and $a_{34} = a_{43} = a_{31} = a_{24} = a_{23} = 0$, then choose $a_{21} = -\alpha_4$,we get that :

$[\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6, \alpha_7, \alpha_8] \longrightarrow [0, \alpha_2 a_{44}^2, 0, 0, a_{44} (\alpha_2 a_{41} + \alpha_5), 0, 1, 0].$

If $\alpha_2 \neq 0$ choose $a_{41} = -\frac{\alpha_5}{\alpha_2}$, so we ge a cocycle $\theta_1 = \delta_{d,d} + \delta_{b,c}$. If $\alpha_2 = 0$ then $\alpha_5 \neq 0$ otherwise $\theta^{\perp} \cap Z(J_{4,2}) \neq 0$, choose $a_{41} = \frac{1}{\alpha_5}$, so we get a cocycle $\theta_2 = \delta_{a,d} + \delta_{b,c}$. A Gröbner basis computation shows that θ_1 and θ_2 are not in the same $Aut(J_{4,2})$-orbit, so we get two algebras :

- $J_{5,7} : a^2 = b, d^2 = e, b \circ c = e.$
- $J_{5,8} : a^2 = b, a \circ d = e, b \circ c = e.$

4.4 1-dimensional centeral extension of $J_{4,3}$

Let $\theta = \alpha_1 \delta_{a,a} + \alpha_2 \delta_{b,b} + \alpha_3 \delta_{d,d} + \alpha_4 \delta_{a,c} + \alpha_5 \delta_{a,d} + \alpha_6 \delta_{b,c} + \alpha_7 \delta_{b,d} + \alpha_8 \delta_{c,d}$ and $(\alpha_4, \alpha_6, \alpha_8) \neq 0$. The center of

$J_{4,3}$, $Z(J_{4,3})$, is spanned by c and d. Furthermore the automorphism group, $Aut(J_{4,3})$, consists of :

$$\phi = \begin{bmatrix} a_{11} & a_{12} & 0 & 0 \\ a_{21} & a_{22} & 0 & 0 \\ a_{31} & a_{32} & a_{11}a_{22} + a_{12}a_{21} & a_{34} \\ a_{41} & a_{42} & 0 & a_{44} \end{bmatrix}, a_{44} \left(a_{11}^2 a_{22}^2 - a_{12}^2 a_{21}^2\right) \neq 0, a_{11}a_{21} = a_{12}a_{22} = 0.$$

The automorphism group acts as follows :

$\alpha_1 \longrightarrow a_{41}\left(\alpha_5 a_{11}+\alpha_3 a_{41}+\alpha_7 a_{21}+\alpha_8 a_{31}\right)+a_{11}\left(\alpha_1 a_{11}+\alpha_4 a_{31}+\alpha_5 a_{41}\right)$
$+a_{21}\left(\alpha_2 a_{21}+\alpha_6 a_{31}+\alpha_7 a_{41}\right)+a_{31}\left(\alpha_4 a_{11}+\alpha_6 a_{21}+\alpha_8 a_{41}\right)$
$\alpha_2 \longrightarrow a_{42}\left(\alpha_5 a_{12}+\alpha_3 a_{42}+\alpha_7 a_{22}+\alpha_8 a_{32}\right)+a_{12}\left(\alpha_1 a_{12}+\alpha_4 a_{32}+\alpha_5 a_{42}\right)$
$+a_{22}\left(\alpha_2 a_{22}+\alpha_6 a_{32}+\alpha_7 a_{42}\right)+a_{32}\left(\alpha_4 a_{12}+\alpha_6 a_{22}+\alpha_8 a_{42}\right)$
$\alpha_3 \longrightarrow a_{44}\left(\alpha_3 a_{44}+2\alpha_8 a_{34}\right)$
$\alpha_4 \longrightarrow \left(a_{11}a_{22}+a_{12}a_{21}\right)\left(\alpha_4 a_{11}+\alpha_6 a_{21}+\alpha_8 a_{41}\right)$
$\alpha_5 \longrightarrow a_{11}\left(\alpha_4 a_{34}+\alpha_5 a_{44}\right)+a_{21}\left(\alpha_6 a_{34}+\alpha_7 a_{44}\right)+a_{41}\left(\alpha_3 a_{44}+\alpha_8 a_{34}\right)+\alpha_8 a_{31}a_{44}$
$\alpha_6 \longrightarrow \left(a_{11}a_{22}+a_{12}a_{21}\right)\left(\alpha_4 a_{12}+\alpha_6 a_{22}+\alpha_8 a_{42}\right)$
$\alpha_7 \longrightarrow a_{12}\left(\alpha_4 a_{34}+\alpha_5 a_{44}\right)+a_{22}\left(\alpha_6 a_{34}+\alpha_7 a_{44}\right)+a_{42}\left(\alpha_3 a_{44}+\alpha_8 a_{34}\right)+\alpha_8 a_{32}a_{44}$
$\alpha_8 \longrightarrow \alpha_8 a_{44}\left(a_{11}a_{22}+a_{12}a_{21}\right)$

Case (1) If $\alpha_8 \neq 0$:

Take $a_{12}=a_{21}=0, a_{11}=a_{22}=a_{44}=1, a_{34}=-\frac{\alpha_3}{2\alpha_8}, a_{41}=-\frac{\alpha_4}{\alpha_8}$ and $a_{42}=-\frac{\alpha_6}{\alpha_8}$ then $[\alpha_3,\alpha_4,\alpha_6] \longrightarrow [0,0,0]$, so we can assume that $\alpha_3=\alpha_4=\alpha_6=0$. Choose $a_{34}=a_{41}=a_{42}=0, a_{31}=-\frac{\alpha_5 a_{11}+\alpha_7 a_{21}}{\alpha_8}$ and $a_{31}=-\frac{\alpha_5 a_{12}+\alpha_7 a_{22}}{\alpha_8}$, it follows that :

$\alpha_1 \longrightarrow \alpha_1 a_{11}^2+\alpha_2 a_{21}^2$,
$\alpha_2 \longrightarrow \alpha_1 a_{12}^2+\alpha_2 a_{22}^2$,
$\alpha_3 \longrightarrow 0, \alpha_4 \longrightarrow 0, \alpha_5 \longrightarrow 0, \alpha_6 \longrightarrow 0, \alpha_7 \longrightarrow 0$,
$\alpha_8 \longrightarrow \alpha_8 a_{44}\left(a_{11}a_{22}+a_{12}a_{21}\right)$.

Case (1.1) If $\alpha_1=\alpha_2=0$: Choose $a_{11}=a_{22}=1$ and $a_{44}=\frac{1}{\alpha_8}$ we get a cocycle $\theta_1=\delta_{c,d}$.

Case (1.2) If $\alpha_1 \neq 0, \alpha_2=0$: Choose $a_{12}=a_{21}=0, a_{11}=\frac{1}{\sqrt{\alpha_1}}, a_{22}=\sqrt{\alpha_1}$ and $a_{44}=\frac{1}{\alpha_8}$ we get a cocycle $\theta_2=\delta_{a,a}+\delta_{c,d}$.

Case (1.3) If $\alpha_1=0, \alpha_2 \neq 0$: Choose $a_{11}=a_{22}=0, a_{12}=\sqrt{\alpha_2}, a_{21}=\frac{1}{\sqrt{\alpha_2}}$ and $a_{44}=\frac{1}{\alpha_8}$ we get again a cocycle $\theta_2=\delta_{a,a}+\delta_{c,d}$.

Case (1.4) If $\alpha_1 \neq 0, \alpha_2 \neq 0$: Choose $a_{12}=a_{21}=0, a_{11}=\frac{1}{\sqrt{\alpha_1}}, a_{22}=\frac{1}{\sqrt{\alpha_2}}$ and $a_{44}=\frac{\sqrt{\alpha_1}\sqrt{\alpha_2}}{\alpha_8}$ we get a cocycle $\theta_3=\delta_{a,a}+\delta_{b,b}+\delta_{c,d}$.

Case (2) If $\alpha_8=0$:

$\alpha_1 \longrightarrow a_{41}\left(\alpha_5 a_{11}+\alpha_3 a_{41}+\alpha_7 a_{21}\right)+a_{11}\left(\alpha_1 a_{11}+\alpha_4 a_{31}+\alpha_5 a_{41}\right)$
$+a_{21}\left(\alpha_2 a_{21}+\alpha_6 a_{31}+\alpha_7 a_{41}\right)+a_{31}\left(\alpha_4 a_{11}+\alpha_6 a_{21}\right)$
$\alpha_2 \longrightarrow a_{42}\left(\alpha_5 a_{12}+\alpha_3 a_{42}+\alpha_7 a_{22}\right)+a_{12}\left(\alpha_1 a_{12}+\alpha_4 a_{32}+\alpha_5 a_{42}\right)$
$+a_{22}\left(\alpha_2 a_{22}+\alpha_6 a_{32}+\alpha_7 a_{42}\right)+a_{32}\left(\alpha_4 a_{12}+\alpha_6 a_{22}\right)$
$\alpha_3 \longrightarrow \alpha_3 a_{44}^2$
$\alpha_4 \longrightarrow \left(a_{11}a_{22}+a_{12}a_{21}\right)\left(\alpha_4 a_{11}+\alpha_6 a_{21}\right)$
$\alpha_5 \longrightarrow a_{11}\left(\alpha_4 a_{34}+\alpha_5 a_{44}\right)+a_{21}\left(\alpha_6 a_{34}+\alpha_7 a_{44}\right)+\alpha_3 a_{44}a_{41}$
$\alpha_6 \longrightarrow \left(a_{11}a_{22}+a_{12}a_{21}\right)\left(\alpha_4 a_{12}+\alpha_6 a_{22}\right)$
$\alpha_7 \longrightarrow a_{12}\left(\alpha_4 a_{34}+\alpha_5 a_{44}\right)+a_{22}\left(\alpha_6 a_{34}+\alpha_7 a_{44}\right)+\alpha_3 a_{44}a_{42}$
$\alpha_8 \longrightarrow 0$

We need at least one of α_4, α_6 to be nonzero in order to have that c do not lie in the radical of θ and at least one of $\alpha_3, \alpha_5, \alpha_7$ to be nonzero in order to have that d do not lie in the radical of θ.

Case (2.1) If $\alpha_3 \neq 0$: We can choose a_{41}, a_{42} and a_{34} such that $[\alpha_5, \alpha_7] \longrightarrow [0,0]$, so we can assume that $\alpha_5 = \alpha_7 = 0$. To fix $[\alpha_5, \alpha_7] \longrightarrow [0,0]$ choose $a_{41} = a_{42} = a_{34} = 0$ then we get that :

$\alpha_1 \longrightarrow \alpha_1 a_{11}^2 + \alpha_2 a_{21}^2 + 2a_{31}(\alpha_4 a_{11} + \alpha_6 a_{21})$,
$\alpha_2 \longrightarrow \alpha_1 a_{12}^2 + \alpha_2 a_{22}^2 + 2a_{32}(\alpha_4 a_{12} + \alpha_6 a_{22})$,
$\alpha_3 \longrightarrow \alpha_3 a_{44}^2$,
$\alpha_4 \longrightarrow (a_{11}a_{22} + a_{12}a_{21})(\alpha_4 a_{11} + \alpha_6 a_{21})$,
$\alpha_6 \longrightarrow (a_{11}a_{22} + a_{12}a_{21})(\alpha_4 a_{12} + \alpha_6 a_{22})$,
$\alpha_5 \longrightarrow 0, \alpha_7 \longrightarrow 0, \alpha_8 \longrightarrow 0$.

Case (2.1.1) If $\alpha_4 \neq 0, \alpha_6 \neq 0$: Choose $a_{12} = a_{21} = 0, a_{11} = \sqrt[3]{\frac{\alpha_6}{\alpha_4^2}}, a_{22} = \sqrt[3]{\frac{\alpha_4}{\alpha_6^2}}, a_{31} = -\frac{\alpha_1}{2\alpha_4}\sqrt[3]{\frac{\alpha_6}{\alpha_4^2}}, a_{32} = -\frac{\alpha_2}{2\alpha_4}\sqrt[3]{\frac{\alpha_4}{\alpha_6^2}}$ and $a_{44} = \frac{1}{\sqrt{\alpha_3}}$ then we get a cocycle $\theta_4 = \delta_{d,d} + \delta_{a,c} + \delta_{b,c}$.

Case (2.1.2) If $\alpha_4 \neq 0, \alpha_6 = 0$: We have that :

$\alpha_1 \longrightarrow \alpha_1 a_{11}^2 + \alpha_2 a_{21}^2 + 2\alpha_4 a_{11} a_{31}$,
$\alpha_2 \longrightarrow \alpha_1 a_{12}^2 + \alpha_2 a_{22}^2 + 2\alpha_4 a_{12} a_{32}$,
$\alpha_3 \longrightarrow \alpha_3 a_{44}^2$,
$\alpha_4 \longrightarrow (a_{11}a_{22} + a_{12}a_{21})\alpha_4 a_{11}$,
$\alpha_6 \longrightarrow (a_{11}a_{22} + a_{12}a_{21})\alpha_4 a_{12}$,
$\alpha_5 \longrightarrow 0, \alpha_7 \longrightarrow 0, \alpha_8 \longrightarrow 0$.

If $\alpha_2 = 0$ choose $a_{12} = a_{21} = 0, a_{11} = \frac{1}{\alpha_4}, a_{22} = 1, a_{31} = -\frac{\alpha_1}{2\alpha_4^2}$ and $a_{44} = \frac{1}{\sqrt{\alpha_3}}$, we get a cocycle $\theta_5 = \delta_{d,d} + \delta_{a,c}$. If $\alpha_2 \neq 0$ choose $a_{12} = a_{21} = 0, a_{11} = \frac{1}{\alpha_4}, a_{22} = \frac{1}{\sqrt{\alpha_2}}, a_{31} = -\frac{\alpha_1}{2\alpha_4^2}$ and $a_{44} = \frac{1}{\sqrt{\alpha_3}}$, we get a cocycle $\theta_6 = \delta_{b,b} + \delta_{d,d} + \delta_{a,c}$.

Case (2.1.3) If $\alpha_4 = 0, \alpha_6 \neq 0$: Choose $a_{11} = a_{22} = 0$, we are back in Case (2.1.2).

Case (2.2) If $\alpha_3 = 0$: We have that :

$\alpha_1 \longrightarrow a_{41}(\alpha_5 a_{11} + \alpha_7 a_{21}) + a_{11}(\alpha_1 a_{11} + \alpha_4 a_{31} + \alpha_5 a_{41})$
$+a_{21}(\alpha_2 a_{21} + \alpha_6 a_{31} + \alpha_7 a_{41}) + a_{31}(\alpha_4 a_{11} + \alpha_6 a_{21})$,
$\alpha_2 \longrightarrow a_{42}(\alpha_5 a_{12} + \alpha_7 a_{22}) + a_{12}(\alpha_1 a_{12} + \alpha_4 a_{32} + \alpha_5 a_{42})$
$+a_{22}(\alpha_2 a_{22} + \alpha_6 a_{32} + \alpha_7 a_{42}) + a_{32}(\alpha_4 a_{12} + \alpha_6 a_{22})$,
$\alpha_4 \longrightarrow (a_{11}a_{22} + a_{12}a_{21})(\alpha_4 a_{11} + \alpha_6 a_{21})$,
$\alpha_5 \longrightarrow a_{11}(\alpha_4 a_{34} + \alpha_5 a_{44}) + a_{21}(\alpha_6 a_{34} + \alpha_7 a_{44})$,
$\alpha_6 \longrightarrow (a_{11}a_{22} + a_{12}a_{21})(\alpha_4 a_{12} + \alpha_6 a_{22})$,
$\alpha_7 \longrightarrow a_{12}(\alpha_4 a_{34} + \alpha_5 a_{44}) + a_{22}(\alpha_6 a_{34} + \alpha_7 a_{44})$,
$\alpha_3 \longrightarrow 0, \alpha_8 \longrightarrow 0$.

Case (2.2.1) If $\alpha_4 \neq 0, \alpha_6 \neq 0$: If we take $a_{12} = a_{21} = 0, a_{34} = -\frac{\alpha_7}{\alpha_6}$ and $a_{44} = 1$ then $\alpha_7 \longrightarrow 0$, so we can assume that $\alpha_7 = 0$. Assume that $\alpha_7 = 0$ then choose $a_{12} =$

$a_{21} = a_{41} = a_{42} = 0, a_{11} = \sqrt[3]{\frac{\alpha_6}{\alpha_4^2}}, a_{22} = \sqrt[3]{\frac{\alpha_4}{\alpha_6^2}}, a_{31} = -\frac{\alpha_1}{2\alpha_4}\sqrt[3]{\frac{\alpha_6}{\alpha_4^2}}, a_{32} = -\frac{\alpha_2}{2\alpha_4}\sqrt[3]{\frac{\alpha_4}{\alpha_6^2}}, a_{34} = 0$ and $a_{44} = \frac{1}{\alpha_5}\sqrt[3]{\frac{\alpha_4^2}{\alpha_6}}$ thus we get a cocycle $\theta_7 = \delta_{a,c} + \delta_{a,d} + \delta_{b,c}$.

Case (2.2.2) If $\alpha_4 \neq 0, \alpha_6 = 0$: Choose $a_{12} = a_{21} = a_{41} = 0, a_{11} = \frac{1}{\sqrt{\alpha_4}}, a_{22} = 1, a_{34} = -\frac{\alpha_5}{\alpha_4\alpha_7}, a_{31} = -\frac{\alpha_1}{2\alpha_4\sqrt{\alpha_4}}, a_{42} = -\frac{\alpha_2}{2\alpha_7}\sqrt[3]{\frac{\alpha_4}{\alpha_6^2}}$ and $a_{44} = \frac{1}{\alpha_7}$ hence we get a cocycle $\theta_8 = \delta_{a,c} + \delta_{b,d}$.

Case (2.2.3) If $\alpha_4 = 0, \alpha_6 \neq 0$: Choose $a_{11} = a_{22} = 0$, we are back in Case (2.2.2).

A Gröbner basis computation shows that the cocycles " $\theta_1, \theta_2, ..., \theta_8$ " are not in the same $Aut(J_{4,3})$-orbit, so we get 8 algebras as follow :

- $J_{5,9} : a \circ b = c, c \circ d = e.$
- $J_{5,10} : a \circ b = c, c \circ d = e, a^2 = e.$
- $J_{5,11} : a \circ b = c, c \circ d = e, a^2 = e, b^2 = e.$
- $J_{5,12} : a \circ b = c, a \circ c = e, d^2 = e, b \circ c = e.$
- $J_{5,13} : a \circ b = c, a \circ c = e, d^2 = e.$
- $J_{5,14} : a \circ b = c, a \circ c = e, d^2 = e, b^2 = e.$
- $J_{5,15} : a \circ b = c, a \circ c = e, a \circ d = e, b \circ c = e.$
- $J_{5,16} : a \circ b = c, a \circ c = e, b \circ d = e.$

4.5 1-dimensional centeral extension of $J_{4,4}$

Let $\theta = \alpha_1 (\delta_{a,c} + \delta_{b,b}) + \alpha_2\delta_{a,d} + \alpha_3\delta_{b,d} + \alpha_4\delta_{d,d}$ such that $\alpha_3 \neq 0$. The center of $J_{4,4}$, $Z(J_{4,4})$, is spanned by c and d. Furthermore the automorphism group, $Aut(J_{4,4})$, consists of :

$$\phi = \begin{bmatrix} a_{11} & 0 & 0 & 0 \\ a_{21} & a_{11}^2 & 0 & 0 \\ a_{31} & 2a_{11}a_{21} & a_{11}^3 & a_{34} \\ a_{41} & 0 & 0 & a_{44} \end{bmatrix}, a_{11}a_{44} \neq 0.$$

The automorphism group acts as follows :

$\alpha_1 \longrightarrow \alpha_1 a_{11}^4,$
$\alpha_2 \longrightarrow a_{44} (\alpha_2 a_{11} + \alpha_3 a_{21} + \alpha_4 a_{41}) + \alpha_1 a_{11} a_{34},$
$\alpha_3 \longrightarrow \alpha_3 a_{11}^2 a_{44},$
$\alpha_4 \longrightarrow \alpha_4 a_{44}^2.$

We need $\alpha_1 \neq 0$ in order to have that c do not lie in the radical of θ. Choose $a_{21} = a_{41} = 0$ and $a_{34} = -\frac{\alpha_2 a_{44}}{\alpha_1}$ to have $\alpha_2 \longrightarrow 0$. So we can assume that $\theta = \alpha_1 (\delta_{a,c} + \delta_{b,b}) + \alpha_3 \delta_{b,d} + \alpha_4 \delta_{d,d}$. To fix $\alpha_2 \longrightarrow 0$ choose $a_{21} = a_{41} = a_{34} = 0$, then :

$\alpha_1 \longrightarrow \alpha_1 a_{11}^4, \alpha_2 \longrightarrow 0, \alpha_3 \longrightarrow \alpha_3 a_{11}^2 a_{44}, \alpha_4 \longrightarrow \alpha_4 a_{44}^2$.

Choose $a_{11}^2 = \frac{1}{\sqrt{\alpha_1}}$ and $a_{44} = \frac{\sqrt{\alpha_1}}{\alpha_3}$, then we get the cocycles $\theta_\alpha = \delta_{a,c} + \delta_{b,b} + \delta_{b,d} + \alpha \delta_{d,d}$. Now the problem is to describe exactly when there exists a $\phi \in Aut(J_{4,4})$ with $\phi\theta_\alpha = \lambda\theta_\beta$ with $\alpha, \beta \in \Bbbk$ and $\lambda \in \Bbbk^*$. A Gröbner basis computation shows that θ_α and θ_β are in the same $Aut(J_{4,4})$-orbit if and only if $\alpha = \beta$ so we get algebras :

- $J_{5,17}^{\alpha} : a^2 = b, a \circ b = c, a \circ c = e, b^2 = e, b \circ d = e, d^2 = \alpha e \quad , \alpha \in \Bbbk$.

4.6 1-dimensional centeral extension of $J_{4,5}$

Firstly before we consider 1-dimensional centeral extension of $J_{4,5}$, we state the following useful lemma which we will use in computation of 1-dimensional centeral extension of $J_{4,5}$.

Lemma 4.1. *Let* $\alpha = [\alpha_1, \alpha_2, \alpha_3]$, $\alpha \neq 0$ *then there exist* $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$ *such that* $\det A \neq 0$ *and*

$$\begin{bmatrix} a_{23} & a_{13} & a_{33} \\ a_{22} & a_{12} & a_{32} \\ a_{21} & a_{11} & a_{31} \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} = \begin{bmatrix} 0 & 0 & c \\ c & 0 & 0 \\ 0 & c & 0 \end{bmatrix} , c \neq 0$$

satisfying $\alpha A = [1, 0, 0]$ *or* $\alpha A = [0, 0, 1]$.

Proof. Case (1) $\alpha_3 = 0$:

Case (1.1) If $\alpha_1 \neq 0, \alpha_2 = 0$ take

$$A = \begin{bmatrix} \frac{1}{\alpha_1} & 0 & 0 \\ 0 & \alpha_1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Case (1.2) If $\alpha_1 = 0, \alpha_2 \neq 0$ take

$$A = \begin{bmatrix} 0 & \alpha_2 & 0 \\ \frac{1}{\alpha_2} & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Case (1.3) If $\alpha_1 \neq 0, \alpha_2 \neq 0$ take

$$A = \begin{bmatrix} -\sqrt{\frac{\alpha_2}{8\alpha_1^3}} & \sqrt{\frac{1}{8\alpha_1\alpha_2}} & \frac{1}{2\alpha_1} \\ \sqrt{\frac{1}{8\alpha_1\alpha_2}} & -\sqrt{\frac{\alpha_1}{8\alpha_2^3}} & \frac{1}{2\alpha_2} \\ \frac{1}{2\alpha_1} & \frac{1}{2\alpha_2} & 0 \end{bmatrix}$$

Case (2) $\alpha_3 \neq 0$:
Case (2.1) If $\alpha_1 \neq 0, \alpha_2 = 0$ take

$$A = \begin{bmatrix} 0 & -\frac{\alpha_3}{\alpha_1} & 0 \\ -\frac{\alpha_1}{\alpha_3^3} & \frac{\alpha_1}{2\alpha_3} & \frac{\alpha_1}{\alpha_3^2} \\ 0 & 1 & \frac{1}{\alpha_3} \end{bmatrix}$$

Case (2.2) If $\alpha_1 = 0, \alpha_2 \neq 0$ take

$$A = \begin{bmatrix} -\frac{\alpha_2}{\alpha_3^3} & \frac{\alpha_2}{2\alpha_3} & \frac{\alpha_2}{\alpha_3^2} \\ 0 & -\frac{\alpha_3}{\alpha_2} & 0 \\ 0 & 1 & \frac{1}{\alpha_3} \end{bmatrix}$$

Case (2.3) If $\alpha_1 \neq 0, \alpha_2 \neq 0$ and $2\alpha_1\alpha_2 + \alpha_3^2 \neq 0$ take

$$A = \begin{bmatrix} \frac{\sqrt{\alpha_1\alpha_2}\left(-\sqrt{2\alpha_1\alpha_2+\alpha_3^2}-\alpha_3\right)}{2\alpha_1\left(2\alpha_1\alpha_2+\alpha_3^2\right)} & \frac{\sqrt{\alpha_1\alpha_2}\left(\sqrt{2\alpha_1\alpha_2+\alpha_3^2}-\alpha_3\right)}{2\alpha_1\left(2\alpha_1\alpha_2+\alpha_3^2\right)} & \frac{\alpha_2}{2\alpha_1\alpha_2+\alpha_3^2} \\ \frac{\sqrt{\alpha_1\alpha_2}\left(\sqrt{2\alpha_1\alpha_2+\alpha_3^2}-\alpha_3\right)}{2\alpha_2\left(2\alpha_1\alpha_2+\alpha_3^2\right)} & \frac{\sqrt{\alpha_1\alpha_2}\left(-\sqrt{2\alpha_1\alpha_2+\alpha_3^2}-\alpha_3\right)}{2\alpha_2\left(2\alpha_1\alpha_2+\alpha_3^2\right)} & \frac{\alpha_1}{2\alpha_1\alpha_2+\alpha_3^2} \\ \frac{\sqrt{\alpha_1\alpha_2}}{2\alpha_1\alpha_2+\alpha_3^2} & \frac{\sqrt{\alpha_1\alpha_2}}{2\alpha_1\alpha_2+\alpha_3^2} & \frac{\alpha_3}{2\alpha_1\alpha_2+\alpha_3^2} \end{bmatrix}$$

Case (2.4) If $\alpha_1 \neq 0, \alpha_2 \neq 0$ and $2\alpha_1\alpha_2 + \alpha_3^2 = 0$ take

$$A = \begin{bmatrix} \frac{1}{4\alpha_1} & \frac{-\alpha_3^2}{2\alpha_1} & \frac{-\alpha_3}{2\alpha_1} \\ \frac{1}{4\alpha_2} & \frac{-\alpha_3^2}{2\alpha_2} & \frac{\alpha_3}{2\alpha_2} \\ \frac{1}{2\alpha_3} & \alpha_3 & 0 \end{bmatrix}$$

□

Now we consider that 1-dimensional centeral extension of $J_{4,5}$. Let $\theta = \alpha_1\delta_{a,d} + \alpha_2\delta_{b,d} + \alpha_3\delta_{c,d} + \alpha_4\delta_{a,a} + \alpha_5\delta_{b,b} + \alpha_6\delta_{c,c} + \alpha_7\delta_{a,b} + \alpha_8\delta_{a,c} + \alpha_9\delta_{b,c}$ modulo $\delta_{c,c} + \delta_{a,b}$ such that $(\alpha_1, \alpha_2, \alpha_3) \neq (0, 0, 0)$. The center of $J_{4,7}$, $Z(J_{4,5})$, is spanned by d. Furthermore the automorphism group, $Aut(J_{4,5})$, consists of :

$$\phi = \begin{bmatrix} a_{11} & a_{12} & a_{13} & 0 \\ a_{21} & a_{22} & a_{23} & 0 \\ a_{31} & a_{32} & a_{33} & 0 \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}, \det\phi \neq 0,$$

Such that :

$$\begin{bmatrix} a_{23} & a_{13} & a_{33} \\ a_{22} & a_{12} & a_{32} \\ a_{21} & a_{11} & a_{31} \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} = \begin{bmatrix} 0 & 0 & a_{44} \\ a_{44} & 0 & 0 \\ 0 & a_{44} & 0 \end{bmatrix}.$$

The automorphism group acts as follows :

$\alpha_1 \longrightarrow a_{44}\left(\alpha_1 a_{11}+\alpha_2 a_{21}+\alpha_3 a_{31}\right)$

$\alpha_2 \longrightarrow a_{44}\left(\alpha_1 a_{12}+\alpha_2 a_{22}+\alpha_3 a_{32}\right)$

$\alpha_3 \longrightarrow a_{44}\left(\alpha_1 a_{13}+\alpha_2 a_{23}+\alpha_3 a_{33}\right)$

$\alpha_4 \longrightarrow \alpha_4 a_{11}^2+2\alpha_7 a_{11}a_{21}+2\alpha_8 a_{11}a_{31}+2\alpha_1 a_{41}a_{11}+\alpha_5 a_{21}^2+2\alpha_9 a_{21}a_{31}$
$+2\alpha_2 a_{41}a_{21}+\alpha_6 a_{31}^2+2\alpha_3 a_{41}a_{31}$

$\alpha_5 \longrightarrow \alpha_4 a_{12}^2+2\alpha_7 a_{12}a_{22}+2\alpha_8 a_{12}a_{32}+2\alpha_1 a_{42}a_{12}+\alpha_5 a_{22}^2+2\alpha_9 a_{22}a_{32}$
$+2\alpha_2 a_{42}a_{22}+\alpha_6 a_{32}^2+2\alpha_3 a_{42}a_{32}$

$\alpha_6 \longrightarrow \alpha_4 a_{13}^2+2\alpha_7 a_{13}a_{23}+2\alpha_8 a_{13}a_{33}+2\alpha_1 a_{43}a_{13}+\alpha_5 a_{23}^2+2\alpha_9 a_{23}a_{33}$
$+2\alpha_2 a_{43}a_{23}+\alpha_6 a_{33}^2+2\alpha_3 a_{43}a_{33}$

$\alpha_7 \longrightarrow a_{12}\left(\alpha_1 a_{41}+\alpha_4 a_{11}+\alpha_7 a_{21}+\alpha_8 a_{31}\right)+a_{22}\left(\alpha_2 a_{41}+\alpha_5 a_{21}+\alpha_7 a_{11}+\alpha_9 a_{31}\right)$
$+a_{32}\left(\alpha_3 a_{41}+\alpha_6 a_{31}+\alpha_8 a_{11}+\alpha_9 a_{21}\right)+a_{42}\left(\alpha_1 a_{11}+\alpha_2 a_{21}+\alpha_3 a_{31}\right)$

$\alpha_8 \longrightarrow a_{13}\left(\alpha_1 a_{41}+\alpha_4 a_{11}+\alpha_7 a_{21}+\alpha_8 a_{31}\right)+a_{23}\left(\alpha_2 a_{41}+\alpha_5 a_{21}+\alpha_7 a_{11}+\alpha_9 a_{31}\right)$
$+a_{33}\left(\alpha_3 a_{41}+\alpha_6 a_{31}+\alpha_8 a_{11}+\alpha_9 a_{21}\right)+a_{43}\left(\alpha_1 a_{11}+\alpha_2 a_{21}+\alpha_3 a_{31}\right)$

$\alpha_9 \longrightarrow a_{13}\left(\alpha_1 a_{42}+\alpha_4 a_{12}+\alpha_7 a_{22}+\alpha_8 a_{32}\right)+a_{23}\left(\alpha_2 a_{42}+\alpha_5 a_{22}+\alpha_7 a_{12}+\alpha_9 a_{32}\right)$
$+a_{33}\left(\alpha_3 a_{42}+\alpha_6 a_{32}+\alpha_8 a_{12}+\alpha_9 a_{22}\right)+a_{43}\left(\alpha_1 a_{12}+\alpha_2 a_{22}+\alpha_3 a_{32}\right)$

By Lemma 4.1 we consider only two cases :

Case (1) : Assume that $[\alpha_1,\alpha_2,\alpha_3]=[1,0,0]$. Choose $a_{12}=a_{13}=0$ then $a_{44}=a_{11}a_{22}, a_{33}^2=a_{44}, a_{32}=0, a_{11}a_{23}+a_{31}a_{33}=0, 2a_{33}a_{21}=a_{31}a_{23}, a_{31}a_{22}+a_{23}a_{33}=0$. It follows that :

$\alpha_1 \longrightarrow a_{44}a_{11}$

$\alpha_2 \longrightarrow 0$

$\alpha_3 \longrightarrow 0$

$\alpha_4 \longrightarrow \alpha_4 a_{11}^2+2\alpha_7 a_{11}a_{21}+2\alpha_8 a_{11}a_{31}+2a_{41}a_{11}+\alpha_5 a_{21}^2+2\alpha_9 a_{21}a_{31}+\alpha_6 a_{31}^2$

$\alpha_5 \longrightarrow \alpha_5 a_{22}^2$

$\alpha_6 \longrightarrow \alpha_5 a_{23}^2+2\alpha_9 a_{23}a_{33}+\alpha_6 a_{33}^2$

$\alpha_7 \longrightarrow a_{22}\left(\alpha_5 a_{21}+\alpha_7 a_{11}+\alpha_9 a_{31}\right)+a_{42}a_{11}$

$\alpha_8 \longrightarrow a_{23}\left(\alpha_5 a_{21}+\alpha_7 a_{11}+\alpha_9 a_{31}\right)+a_{33}\left(\alpha_6 a_{31}+\alpha_8 a_{11}+\alpha_9 a_{21}\right)+a_{43}a_{11}$

$\alpha_9 \longrightarrow a_{22}\left(a_{23}\alpha_5+a_{33}\alpha_9\right).$

Case (1.1) : If $\alpha_5=0$ choose $a_{23}=0$ then $a_{44}=a_{11}a_{22}, a_{33}^2=a_{44}, a_{32}=0, a_{31}=0, a_{21}=0$. It follows that :

$\alpha_1 \longrightarrow a_{44}a_{11}$, $\alpha_2 \longrightarrow 0$, $\alpha_3 \longrightarrow 0$,

$\alpha_4 \longrightarrow a_{11}\left(\alpha_4 a_{11}+2a_{41}\right)$, $\alpha_5 \longrightarrow 0$, $\alpha_6 \longrightarrow \alpha_6 a_{33}^2$,

$\alpha_7 \longrightarrow a_{11}\left(\alpha_7 a_{22}+a_{42}\right)$, $\alpha_8 \longrightarrow a_{11}\left(\alpha_8 a_{33}+a_{43}\right)$,

$\alpha_9 \longrightarrow \alpha_9 a_{22}a_{33}$.

Case (1.1.1) : If $\alpha_9=0$ choose $a_{11}=a_{22}=1$ " $\Rightarrow a_{33}=a_{44}=1$", $a_{41}=\frac{-\alpha_4}{2}, a_{42}=\alpha_6-\alpha_7$ and $a_{43}=-\alpha_8$, we get a cocycle $\theta_1=\delta_{a,d}$.

Case (1.1.2) : If $\alpha_9\neq 0$ choose $a_{11}=\alpha_9^{\frac{2}{5}}, a_{22}=\alpha_9^{\frac{-4}{5}}$ " $\Rightarrow a_{33}^2=a_{44}=\alpha_9^{\frac{-2}{5}}$", $a_{41}=\frac{-\alpha_4}{2}, a_{42}=\alpha_9^{\frac{-2}{5}}\alpha_6-\alpha_9^{\frac{-4}{5}}\alpha_7$ and $a_{43}=-\alpha_8$, we get a cocycle $\theta_2=\delta_{a,d}+\delta_{b,c}$.

Case (1.2) : If $\alpha_5 \neq 0$ choose $a_{23} = \frac{-a_{33}\alpha_9}{\alpha_5}$ then $\alpha_9 \longrightarrow 0$, thus we can assume that $\alpha_9 = 0$ and choose $a_{23} = 0$ to keep $\alpha_9 \longrightarrow 0$. It follows that :

$\alpha_1 \longrightarrow a_{44}a_{11}, \alpha_2 \longrightarrow 0, \alpha_3 \longrightarrow 0, \alpha_4 \longrightarrow a_{11}(\alpha_4 a_{11} + 2a_{41}), \alpha_5 \longrightarrow \alpha_5 a_{22}^2,$

$\alpha_6 \longrightarrow \alpha_6 a_{33}^2, \alpha_7 \longrightarrow a_{11}(\alpha_7 a_{22} + a_{42}), \alpha_8 \longrightarrow a_{11}(\alpha_8 a_{33} + a_{43}), \alpha_9 \longrightarrow 0.$

Choose $a_{11} = \sqrt[4]{\alpha_5}, a_{22} = \frac{1}{\sqrt{\alpha_5}}$ $\left(\text{then } a_{33}^2 = a_{44} = \frac{1}{\sqrt[4]{\alpha_5}}\right), a_{41} = \frac{-\alpha_4}{2}, a_{42} = \frac{1}{\sqrt[4]{\alpha_5}}\alpha_6 - \frac{1}{\sqrt{\alpha_5}}\alpha_7$ and $a_{43} = -\alpha_8$, we get a cocycle $\theta_3 = \delta_{a,d} + \delta_{b,b}$.

Case (2) : Assume that $[\alpha_1, \alpha_2, \alpha_3] = [0, 0, 1]$. Choose $a_{31} = a_{32} = a_{13} = a_{23} = a_{12} = a_{21} = 0, a_{11} = a_{22} = a_{33} = a_{44} = 1$ it follows that :

$\alpha_1 \longrightarrow 0, \alpha_2 \longrightarrow 0, \alpha_3 \longrightarrow 0, \alpha_4 \longrightarrow \alpha_4, \alpha_5 \longrightarrow \alpha_5,$

$\alpha_6 \longrightarrow \alpha_6 + 2a_{43}, \alpha_7 \longrightarrow \alpha_7 + a_{42}, \alpha_8 \longrightarrow a_{41} + \alpha_8,$

$\alpha_9 \longrightarrow a_{42} + \alpha_9.$

Choose $a_{41} = -\alpha_8, a_{42} = -\alpha_9, a_{43} = \frac{\alpha_7 - \alpha_9 - 2\alpha_6}{2}$ we get a cocycles $\theta_4^{\alpha,\beta} = \delta_{c,d} + \alpha\delta_{a,a} + \beta\delta_{b,b}$. So we get the algebra $J_{\theta_4^{\alpha,\beta}} : a \circ b = d, c^2 = d, c \circ d = e, a^2 = \alpha e, b^2 = \beta e$.

Case (2.1) : If $\alpha = \beta = 0$, we get a cocycle $\theta_4^{0,0} = \delta_{c,d}$. Case (2.2) If $\alpha \neq 0, \beta = 0$, putting $a' = \frac{1}{\sqrt{\alpha}}a, b' = \sqrt{\alpha}b, c' = c, d' = d, e' = e$ we get $J_{\theta_4^{\alpha,\beta}} \approxeq J_{\theta_4^{1,0}}$. Then $\theta_4^{\alpha,0}$ and $\theta_4^{1,0}$ are in the same $Aut(J_{4,5})$-orbit, so we may assume $\alpha = 1$. Case (2.3) If $\alpha = 0, \beta \neq 0$, putting $a' = \frac{1}{\sqrt{\beta}}b, b' = \sqrt{\beta}a, c' = c, d' = d, e' = e$ we get $J_{\theta_4^{0,\beta}} \approxeq J_{\theta_4^{1,0}}$. Then $\theta_4^{0,\beta}$ and $\theta_4^{1,0}$ are in the same $Aut(J_{4,5})$-orbit, so we are back in case (2.1). Case (2.4) If $\alpha \neq 0, \beta \neq 0$, putting $a' = \frac{\sqrt[4]{\alpha^5\beta^3}}{\alpha}a, b' = \sqrt[4]{\alpha^3\beta}b, c' = \sqrt{\alpha\beta}c, d' = \alpha\beta d, e' = \alpha\beta\sqrt{\alpha\beta}e$ we get the same multiplication table, but the parameters α, β have changed to $\alpha = \beta = 1$, i.e $J_{\theta_4^{\alpha,\beta}} \approxeq J_{\theta_4^{1,1}}$. Then $\theta_4^{\alpha,\beta}$ and $\theta_4^{1,1}$ are in the same $Aut(J_{4,5})$-orbit, so we may assume $\alpha = \beta = 1$. A Gröbner basis computation shows that the cocycles $\theta_1 = \delta_{a,d}, \theta_2 = \delta_{a,d} + \delta_{b,c}, \theta_3 = \delta_{a,d} + \delta_{b,b}, \theta_4^{0,0} = \delta_{c,d}, \theta_4^{1,0} = \delta_{c,d} + \delta_{a,a}$ and $\theta_4^{1,1} = \delta_{c,d} + \delta_{a,a} + \delta_{b,b}$ are not in the same $Aut(J_{4,5})$-orbit. So we get algebras :

- $J_{5,18} : a \circ b = d, c^2 = d, a \circ d = e.$
- $J_{5,19} : a \circ b = d, c^2 = d, a \circ d = e, b \circ c = e.$
- $J_{5,20} : a \circ b = d, c^2 = d, a \circ d = e, b^2 = e.$
- $J_{5,21} : a \circ b = d, c^2 = d, c \circ d = e.$
- $J_{5,22} : a \circ b = d, c^2 = d, c \circ d = e, a^2 = e.$
- $J_{5,23} : a \circ b = d, c^2 = d, c \circ d = e, a^2 = e, b^2 = e.$

4.7 1-dimensional centeral extension of $J_{4,7}$

Let $\theta = \alpha_1 (\delta_{b,b} + \delta_{a,d}) + \alpha_2\delta_{a,c} + \alpha_3\delta_{a,b} + \alpha_4\delta_{c,c}$ modulo $\delta_{a,b} + \delta_{c,c}$ such that $\alpha_1 \neq 0$.The centre of $J_{4,7}$, $Z(J_{4,7})$, is spanned by d. Furthermore the automorphism group, $Aut(J_{4,7})$, consists of :

$$\phi = \begin{bmatrix} a_{11} & 0 & 0 & 0 \\ a_{21} & a_{11}^2 & -a_{11}^{\frac{1}{2}}a_{31} & 0 \\ a_{31} & 0 & a_{11}^{\frac{3}{2}} & 0 \\ a_{41} & 2a_{11}a_{21} + a_{31} & a_{43} & a_{11}^3 \end{bmatrix}, a_{11} \neq 0.$$

The automorphism group acts as follows :
$\alpha_1 \longrightarrow \alpha_1 a_{11}^4$
$\alpha_2 \longrightarrow \sqrt{a_{11}} \left(\alpha_2 a_{11}^2 + \alpha_1\sqrt{a_{11}}a_{43} - \alpha_1 a_{21}a_{31} - \alpha_3 a_{11}a_{31} + \alpha_4 a_{11}a_{31}\right)$
$\alpha_3 \longrightarrow a_{11} \left(\alpha_3 a_{11}^2 + 3\alpha_1 a_{21}a_{11} + \alpha_1 a_{31}\right)$
$\alpha_4 \longrightarrow \alpha_4 a_{11}^3 + \alpha_1 a_{11}a_{31}^2$
After dividing we can assume that $\alpha_1 = 1$. Now take $a_{11} = 1, a_{21} = 0, a_{31} = \frac{1+\sqrt{1-4(\alpha_4-\alpha_3)}}{2}$ and $a_{43} = -\alpha_2 + (\alpha_3 - \alpha_4)\frac{1+\sqrt{1-4(\alpha_4-\alpha_3)}}{2}$, we obtain a cocycle $\theta = \delta_{b,b} + \delta_{a,d}$. So we get the algebra :

- $J_{5,24} : a^2 = b, a \circ b = d, c^2 = d, b^2 = e, a \circ d = e.$

4.8 1-dimensional centeral extension of $J_{4,12}$

Let $\theta = \alpha_1\delta_{b,b} + \alpha_2\delta_{a,c} + \alpha_3\delta_{b,c} + \alpha_4\delta_{a,d} + \alpha_5\delta_{b,d}$ such that$\alpha_5 \neq 0$ or $\alpha_3 \neq \alpha_4$. The center of $J_{4,12}$, $Z(J_{4,12})$, is spanned by c and d. Furthermore the automorphism group, $Aut(J_{4,12})$, consists of :

$$\phi = \begin{bmatrix} a_{11} & 0 & 0 & 0 \\ a_{21} & a_{22} & 0 & 0 \\ a_{31} & a_{32} & a_{11}^2 & 0 \\ a_{41} & a_{42} & 2a_{11}a_{21} & a_{11}a_{22} \end{bmatrix}, a_{11}a_{22} \neq 0.$$

The automorphism group acts as follows :
$\alpha_1 \longrightarrow a_{22} \left(\alpha_1 a_{22} + 2\alpha_3 a_{32} + 2\alpha_5 a_{42}\right)$
$\alpha_2 \longrightarrow a_{11}^2 \left(\alpha_2 a_{11} + \alpha_3 a_{21}\right) + 2a_{11}a_{21} \left(\alpha_4 a_{11} + \alpha_5 a_{21}\right)$
$\alpha_3 \longrightarrow a_{22} \left(\alpha_3 a_{11}^2 + 2\alpha_5 a_{21}a_{11}\right)$
$\alpha_4 \longrightarrow a_{11}a_{22} \left(\alpha_4 a_{11} + \alpha_5 a_{21}\right)$
$\alpha_5 \longrightarrow \alpha_5 a_{11}a_{22}^2$
Case (1) if $\alpha_5 \neq 0$:
If we take $a_{21} = -\frac{\alpha_4 a_{11}}{\alpha_5}, a_{32} = 0$ and $a_{42} = -\frac{\alpha_1 a_{22}}{2\alpha_5}$ we get that $\alpha_1 \longrightarrow 0, \alpha_4 \longrightarrow 0$. Thus we can assume that $\alpha_1 = \alpha_4 = 0$ then choose $a_{21} = a_{32} = a_{42} = 0$ we obtain that :

$\alpha_1 \longrightarrow 0$
$\alpha_2 \longrightarrow \alpha_2 a_{11}^3$
$\alpha_3 \longrightarrow \alpha_3 a_{11}^2 a_{22}$
$\alpha_4 \longrightarrow 0$
$\alpha_5 \longrightarrow \alpha_5 a_{11} a_{22}^2$

We need at least one of α_2, α_3 nonzero in order to have that c do not lie in the radical of θ, if $\alpha_2 = 0$ we get a cocycle $\theta_1 = \delta_{b,c} + \delta_{b,d}$. If $\alpha_2 \neq 0$ we get cocycles $\theta_2^\alpha = \delta_{a,c} + \alpha\delta_{b,c} + \delta_{b,d}$.

Case (2) if $\alpha_5 = 0$:
$\alpha_1 \longrightarrow a_{22}\left(\alpha_1 a_{22} + 2\alpha_3 a_{32}\right)$
$\alpha_2 \longrightarrow a_{11}^2\left(\alpha_2 a_{11} + a_{21}\left(\alpha_3 + 2\alpha_4\right)\right)$
$\alpha_3 \longrightarrow \alpha_3 a_{11}^2 a_{22}$
$\alpha_4 \longrightarrow \alpha_4 a_{11}^2 a_{22}$
$\alpha_5 \longrightarrow 0$

If $\alpha_3 + 2\alpha_4 \neq 0$ we can solve for a_{21} to have $\alpha_2 \longrightarrow 0$ then $\alpha_3 \neq 0$, also solve for a_{32} to have $\alpha_1 \longrightarrow 0$. We obtain cocycles $\theta_3^{\alpha\neq 0,1} = \delta_{b,c} + \alpha\delta_{a,d}$. If $\alpha_3 + 2\alpha_4 = 0$ then both of α_3, α_4 are nonzero, solve for a_{32} to have $\alpha_1 \longrightarrow 0$. It follows that :
$\alpha_1 \longrightarrow 0$
$\alpha_2 \longrightarrow \alpha_2 a_{11}^3$
$\alpha_3 \longrightarrow -2\alpha_4 a_{11}^2 a_{22}$
$\alpha_4 \longrightarrow \alpha_4 a_{11}^2 a_{22}$
$\alpha_5 \longrightarrow 0$

If $\alpha_2 \neq 0$, take $a_{11} = \frac{1}{\sqrt[3]{\alpha_2}}$ and $a_{22} = \frac{\sqrt[3]{\alpha_2^2}}{\alpha_4}$ we get a cocycle $\theta_4 = \delta_{a,c} - 2\delta_{b,c} + \delta_{a,d}$. If $\alpha_2 = 0$, take $a_{11} = 1, a_{22} = \frac{-1}{2\alpha_4}$ we get a cocycle $\theta_3^{\alpha=\frac{-1}{2}} = \delta_{b,c} + \frac{-1}{2}\delta_{a,d}$. A Gröbner basis computation shows that $\theta_2^\alpha \sim \theta_2^\beta$ if and only if $\alpha^2 - \beta^2 = 0$ and $\theta_3^{\alpha\neq 0,1} \sim \theta_3^{\beta\neq 0,1}$ if and only if $\alpha = \beta$. Then the cocycles $\theta_1 = \delta_{b,c} + \delta_{b,d}, \theta_2^{\alpha\in\Bbbk} = \delta_{a,c} + \alpha\delta_{b,c} + \delta_{b,d}$, $\theta_3^{\alpha\neq 0,1} = \delta_{b,c} + \alpha\delta_{a,d}$ and $\theta_4 = \delta_{a,c} - 2\delta_{b,c} + \delta_{a,d}$ are not in the same $Aut(J_{4,12})$-orbit. So we get algebras :

- $J_{5,25} : a^2 = c, a \circ b = d, b \circ c = e, b \circ d = e.$
- $J_{5,26}^\alpha : a^2 = c, a \circ b = d, a \circ c = e, b \circ d = e, b \circ c = \alpha e.$
- $J_{5,27}^{\alpha\neq 0,1} : a^2 = c, a \circ b = d, b \circ c = e, a \circ d = \alpha e.$
- $J_{5,28} : a^2 = c, a \circ b = d, a \circ c = e, b \circ c = -2e, a \circ d = e.$

Remark 4.2. $J_{5,26}^\alpha$ is isomorphic to $J_{5,26}^\beta$ if and only if $\beta = \alpha$ or $\beta = -\alpha$. To describe the isomorphism, let a, b, c, d, e be the basis elements of $J_{5,26}^\alpha$ and a', b', c', d', e' those of $J_{5,26}^\beta$. Then $\varphi(a) = a', \varphi(b) = -b', \varphi(c) = c', \varphi(d) = -d', \varphi(e) = e'$, defines an isomorphism $\varphi : J_{5,26}^\alpha \longrightarrow J_{5,26}^{-\alpha}$.

4.9 1-dimensional centeral extension of $J_{4,13}$

Let $\theta = \alpha_1\delta_{a,b} + \alpha_2\delta_{a,c} + \alpha_3\delta_{b,d} + \alpha_4\delta_{b,c} + \alpha_5\delta_{a,d}$ such that $(\alpha_4, \alpha_5) \neq (0,0)$. The center of $J_{4,13}$, $Z(J_{4,13})$, is spanned by c and d. Furthermore the automorphism group, $Aut(J_{4,13})$, consists of :

$$\phi = \begin{bmatrix} a_{11} & a_{12} & 0 & 0 \\ a_{21} & a_{22} & 0 & 0 \\ a_{31} & a_{32} & a_{11}^2 & a_{12}^2 \\ a_{41} & a_{42} & a_{21}^2 & a_{22}^2 \end{bmatrix}, a_{11}^3a_{22}^3 - a_{12}^3a_{21}^3 \neq 0, a_{11}a_{12} = a_{22}a_{21} = 0.$$

The automorphism group acts as follows :

$\alpha_1 \longrightarrow a_{11}(\alpha_1a_{22} + \alpha_2a_{32} + \alpha_5a_{42}) + a_{21}(\alpha_1a_{12} + \alpha_3a_{42} + \alpha_4a_{32}) + a_{31}(\alpha_2a_{12} + \alpha_4a_{22}) + a_{41}(\alpha_3a_{22} + \alpha_5a_{12})$

$\alpha_2 \longrightarrow a_{11}^2(\alpha_2a_{11} + \alpha_4a_{21}) + a_{21}^2(\alpha_3a_{21} + \alpha_5a_{11})$

$\alpha_3 \longrightarrow a_{12}^2(\alpha_2a_{12} + \alpha_4a_{22}) + a_{22}^2(\alpha_3a_{22} + \alpha_5a_{12})$

$\alpha_4 \longrightarrow a_{11}^2(\alpha_2a_{12} + \alpha_4a_{22}) + a_{21}^2(\alpha_3a_{22} + \alpha_5a_{12})$

$\alpha_5 \longrightarrow a_{12}^2(\alpha_2a_{11} + \alpha_4a_{21}) + a_{22}^2(\alpha_3a_{21} + \alpha_5a_{11})$

Since $(\alpha_4, \alpha_5) \neq (0,0)$ we can choose a_{31}, a_{32}, a_{41} and a_{42} such that $\alpha_1 \longrightarrow 0$. So we assume that $\alpha_1 = 0$ then choose $a_{31} = a_{32} = a_{41} = a_{42} = 0$ to fix $\alpha_1 \longrightarrow 0$. Then we get that :

$\alpha_1 \longrightarrow 0$

$\alpha_2 \longrightarrow a_{11}^2(\alpha_2a_{11} + \alpha_4a_{21}) + a_{21}^2(\alpha_3a_{21} + \alpha_5a_{11})$

$\alpha_3 \longrightarrow a_{12}^2(\alpha_2a_{12} + \alpha_4a_{22}) + a_{22}^2(\alpha_3a_{22} + \alpha_5a_{12})$

$\alpha_4 \longrightarrow a_{11}^2(\alpha_2a_{12} + \alpha_4a_{22}) + a_{21}^2(\alpha_3a_{22} + \alpha_5a_{12})$

$\alpha_5 \longrightarrow a_{12}^2(\alpha_2a_{11} + \alpha_4a_{21}) + a_{22}^2(\alpha_3a_{21} + \alpha_5a_{11})$

Case (1) If $\alpha_4 = 0, \alpha_5 \neq 0$:

Choose $a_{12} = a_{21} = 0$ it follows that :

$\alpha_1 \longrightarrow 0$, $\alpha_2 \longrightarrow \alpha_2a_{11}^3$, $\alpha_3 \longrightarrow \alpha_3a_{22}^3$, $\alpha_4 \longrightarrow \alpha_4a_{11}^2a_{22}$, $\alpha_5 \longrightarrow \alpha_5a_{11}a_{22}^2$.

We need α_2, α_5 are nonzeros in order to have that c and d do not lie in the radical of θ, so after dividing we get cocycles $\theta_1^\alpha = \delta_{a,c} + \delta_{a,d} + \alpha\delta_{b,d}$.

Case (2) If $\alpha_4 = 0, \alpha_5 \neq 0$:

Choose $a_{11} = a_{22} = 0$ then we are back in case (1).

Case (3) If $\alpha_4 \neq 0, \alpha_5 \neq 0$:

After dividing we get cocycles $\theta_2^{\alpha,\beta} = \delta_{b,c} + \delta_{a,d} + \alpha\delta_{b,d} + \beta\delta_{a,c}$.

A Gröbner basis computation shows that $\theta_1^\alpha \sim \theta_1^{\alpha'}$ if and only if $\alpha = \alpha'$ or $\alpha = -\alpha'$ and $\theta_2^{\alpha,\beta} \sim \theta_2^{\alpha',\beta'}$ if and only if $(\alpha, \beta) = (\alpha', \beta')$ or $(\alpha, \beta) = (\beta', \alpha')$. So we get algebras :

- $J_{5,29}^{\alpha\in\mathbb{k}} : a^2 = c, b^2 = d, a \circ c = e, a \circ d = e, b \circ d = \alpha e.$
- $J_{5,30}^{\alpha,\beta\in\mathbb{k}} : a^2 = c, b^2 = d, b \circ c = e, a \circ d = e, b \circ d = \alpha e, a \circ c = \beta e.$

Remark 4.3. $J^{\alpha}_{5,29}$ is isomorphic to $J^{\beta}_{5,29}$ if and only if $\beta = \alpha$ or $\beta = -\alpha$. In the latter case, $\varphi(a) = a', \varphi(b) = -b', \varphi(c) = c', \varphi(d) = d', \varphi(e) = e'$, defines an isomorphism $\varphi : J^{\alpha}_{5,29} \longrightarrow J^{-\alpha}_{5,29}$.

Remark 4.4. $J^{\alpha,\beta\in\Bbbk}_{5,30}$ is isomorphic to $J^{\alpha',\beta'\in\Bbbk}_{5,30}$ if and only if $(\alpha,\beta) = (\alpha',\beta')$ or $(\alpha,\beta) = (\beta',\alpha')$. In the latter case, $\varphi(a) = b', \varphi(b) = a', \varphi(c) = d', \varphi(d) = c', \varphi(e) = e'$, defines an isomorphism $\varphi : J^{\alpha,\beta\in\Bbbk}_{5,30} \longrightarrow J^{\beta,\alpha\in\Bbbk}_{5,30}$.

4.10 2-dimensional centeral extension of $J_{3,2}$

Here we find the 2-dimensional central extensions of $J_{3,2}$. Let $\theta_1 = \alpha_1\delta_{a,b}+\alpha_2\delta_{a,c}+\alpha_3\delta_{b,c}+\alpha_4\delta_{c,c}$ and $\theta_2 = \alpha_1\delta_{a,b} + \alpha_2\delta_{a,c} + \alpha_3\delta_{b,c} + \alpha_4\delta_{c,c}$ are the basis of a 2-dimensional subspace V of $H^2_{Jor.}(J_{3,2},\Bbbk)$ such that $(\alpha_3,\alpha'_3) \neq (0,0)$. Up to a scalar, we can always identify a 2-dimensional subspace V with the wedge product of θ_1 and θ_2. The center of $J_{3,2}, Z(J_{3,2})$, is spanned by b and c. Furthermore the automorphism group,$Aut(J_{3,2})$, consists of :

$$\phi = \begin{bmatrix} a_{11} & 0 & 0 \\ a_{21} & a_{11}^2 & a_{23} \\ a_{31} & 0 & a_{33} \end{bmatrix}, \; a_{11}a_{33} \neq 0.$$

We need to classify the orbits of this group on a 2-dimensional subspace$V$. The automorphism group acts as follows :

$$\phi \begin{bmatrix} \alpha_1 \\ \alpha_2 \\ \alpha_3 \\ \alpha_4 \end{bmatrix} = \begin{bmatrix} a_{11}^3\alpha_1 + a_{31}a_{11}^2\alpha_3 \\ a_{11}a_{23}\alpha_1 + a_{11}a_{33}\alpha_2 + (a_{21}a_{33} + a_{31}a_{23})\alpha_3 + a_{31}a_{33}\alpha_4 \\ a_{11}^2a_{33}\alpha_3 \\ 2a_{23}a_{33}\alpha_3 + a_{33}^2\alpha_4 \end{bmatrix},$$

$$\phi \begin{bmatrix} \alpha'_1 \\ \alpha'_2 \\ \alpha'_3 \\ \alpha'_4 \end{bmatrix} = \begin{bmatrix} a_{11}^3\alpha'_1 + a_{31}a_{11}^2\alpha'_3 \\ a_{11}a_{23}\alpha'_1 + a_{11}a_{33}\alpha'_2 + (a_{21}a_{33} + a_{31}a_{23})\alpha'_3 + a_{31}a_{33}\alpha'_4 \\ a_{11}^2a_{33}\alpha'_3 \\ 2a_{23}a_{33}\alpha'_3 + a_{33}^2\alpha'_4 \end{bmatrix}.$$

As we require that $(\alpha_3,\alpha'_3) \neq (0,0)$, we may assume $\alpha_3 = 1$ which is always doable " If $\alpha_3 = 0$ then $\alpha'_3 \neq 0$, switch θ_1 and θ_2 so $\alpha_3 \neq 0$ in θ_1, then multiply θ_1 by $\frac{1}{\alpha'_3}$ to get $\alpha_3 = 1$ ". Bear in mind that we can multiply any of our vectors θ_1 and θ_2 by a scalar, as we are dealing essentially with the subspaces, instead of the vectors.

Choose $a_{11} = a_{33} = 1, a_{31} = -\alpha_1, a_{23} = -\frac{\alpha_4}{2}$ and $a_{21} = \alpha_1\alpha_4 - \alpha_2$. It follows that $\phi\theta_1 = \delta_{b,c}$, so we can assume that $\theta_1 = \delta_{b,c}$. Once we get $\theta_1 = \delta_{b,c}$, we may assume $\alpha'_3 = 0$ because we can replace θ_2 by a linear combination $\theta_2 - \theta_1$. Now to fix θ_1 (up to a scalar), we require $a_{31} = a_{23} = a_{21} = 0$.

For θ_2, we have $\alpha'_1 \longrightarrow a_{11}^3\alpha'_1, \alpha'_2 \longrightarrow a_{11}a_{33}\alpha'_2, \alpha'_3 \longrightarrow 0, \alpha'_4 \longrightarrow a_{33}^2\alpha'_4$.

Case (1) If $\alpha_1' \neq 0, \alpha_2' = 0, \alpha_4' = 0$: After dividing we can assume that $\theta_2 = \delta_{a,b}$, then we get a subspace $V_1 = \delta_{b,c} \wedge \delta_{a,b}$.

Case (2) If $\alpha_1' \neq 0, \alpha_2' \neq 0, \alpha_4' = 0$: Choose $a_{11} = \frac{1}{\sqrt[3]{\alpha_1'}}, a_{33} = \frac{\sqrt[3]{\alpha_1'}}{\alpha_2'}$ this lead to $\alpha_1' \longrightarrow 1, \alpha_2' \longrightarrow 1, \alpha_3' \longrightarrow 0, \alpha_4' \longrightarrow 0$, so we can assume that $\theta_2 = \delta_{a,b} + \delta_{a,c}$. Thus we get a subspace $V_2 = \delta_{b,c} \wedge (\delta_{a,b} + \delta_{a,c})$.

Case (3) If $\alpha_1' \neq 0, \alpha_2' = 0, \alpha_4' \neq 0$: Choose $a_{11} = \frac{1}{\sqrt[3]{\alpha_1'}}, a_{33} = \frac{1}{\sqrt{\alpha_4'}}$ this lead to $\alpha_1' \longrightarrow 1, \alpha_2' \longrightarrow 0, \alpha_3' \longrightarrow 0, \alpha_4' \longrightarrow 1$, so we can assume that $\theta_2 = \delta_{a,b} + \delta_{c,c}$. Thus we get a subspace $V_3 = \delta_{b,c} \wedge (\delta_{a,b} + \delta_{c,c})$.

Case (4) If $\alpha_1' = 0, \alpha_2' = 0, \alpha_4' \neq 0$: After dividing we can assume that $\theta_2 = \delta_{c,c}$, then we get a subspace $V_4 = \delta_{b,c} \wedge \delta_{c,c}$.

Case (5) If $\alpha_1' = 0, \alpha_2' \neq 0, \alpha_4' = 0$: After dividing we can assume that $\theta_2 = \delta_{a,c}$, then we get a subspace $V_5 = \delta_{b,c} \wedge \delta_{a,c}$.

Case (6) If $\alpha_1' = 0, \alpha_2' \neq 0, \alpha_4' \neq 0$: Choose $a_{11} = \frac{\sqrt{\alpha_4'}}{\alpha_2'}, a_{33} = \frac{1}{\sqrt{\alpha_4'}}$ this lead to $\alpha_1' \longrightarrow 0, \alpha_2' \longrightarrow 1, \alpha_3' \longrightarrow 0, \alpha_4' \longrightarrow 1$, so we can assume that $\theta_2 = \delta_{a,c} + \delta_{c,c}$. Thus we get a subspace $V_6 = \delta_{b,c} \wedge (\delta_{a,c} + \delta_{c,c})$.

Case (7) If $\alpha_1' \neq 0, \alpha_2' \neq 0, \alpha_4' \neq 0$: Choose $a_{11} = \frac{\alpha_2'^2}{\alpha_1' \alpha_4'}, a_{33} = \frac{\alpha_2'^3}{\alpha_1' \alpha_4'^2}$ this lead to $\alpha_1' \longrightarrow \frac{\alpha_2'^6}{\alpha_1'^2 \alpha_4'^3}, \alpha_2' \longrightarrow \frac{\alpha_2'^6}{\alpha_1'^2 \alpha_4'^3}, \alpha_3' \longrightarrow 0, \alpha_4' \longrightarrow \frac{\alpha_2'^6}{\alpha_1'^2 \alpha_4'^3}$, so after dividing we can assume that $\theta_2 = \delta_{a,b} + \delta_{a,c} + \delta_{c,c}$. Thus we get a subspace $V_7 = \delta_{b,c} \wedge (\delta_{a,b} + \delta_{a,c} + \delta_{c,c})$.

A Gröbner basis computation shows $J_{V_3} : a^2 = b, b \circ c = d, a \circ b = e, c^2 = e$ is isomorphic to $J_{V_7} : a^2 = b, b \circ c = d, a \circ b = e, a \circ c = e, c^2 = e$ by a map $a \longrightarrow a + \frac{4}{9}b + \frac{2}{3}c$, $b \longrightarrow b + \frac{16}{27}d + \frac{4}{3}e, c \longrightarrow \frac{1}{3}b + c, d \longrightarrow d, e \longrightarrow \frac{2}{3}d + e$. Also by Gröbner basis computation we get that$V_1, V_2, ..., V_6$ are not in the same $Aut(J_{3,2})$-orbit. So we get only 6 algebras as follow :

- $J_{5,31} : a^2 = b, b \circ c = d, a \circ b = e.$
- $J_{5,32} : a^2 = b, b \circ c = d, a \circ b = e, a \circ c = e.$
- $J_{5,33} : a^2 = b, b \circ c = d, a \circ b = e, c^2 = e.$
- $J_{5,34} : a^2 = b, b \circ c = d, c^2 = e.$
- $J_{5,35} : a^2 = b, b \circ c = d, a \circ c = e.$
- $J_{5,36} : a^2 = b, b \circ c = d, a \circ c = e, c^2 = e.$

4.11 2-dimensional centeral extension of $J_{3,3}$

Let $\theta_1 = \alpha_1\delta_{a,a} + \alpha_2\delta_{b,b} + \alpha_3\delta_{a,c} + \alpha_4\delta_{b,c}$ and $\theta_2 = \alpha'_1\delta_{a,a} + \alpha'_2\delta_{b,b} + \alpha'_3\delta_{a,c} + \alpha'_4\delta_{b,c}$ are the basis of a 2-dimensional subspace V of $H^2_{Jor.}(J_{3,3}, \Bbbk)$. The center of $J_{3,3}, Z(J_{3,3})$, is spanned by c. Furthermore the automorphism group,$Aut(J_{3,3})$, consists of :

$$\phi = \begin{bmatrix} a_{11} & a_{12} & 0 \\ a_{21} & a_{22} & 0 \\ a_{31} & a_{32} & a_{11}a_{22} + a_{21}a_{12} \end{bmatrix}, a_{11}^2a_{22}^2 - a_{12}^2a_{21}^2 \neq 0 \text{ and } \ a_{11}a_{21} = a_{12}a_{22} = 0$$

The automorphism group acts as follows :

$$\phi \begin{bmatrix} \alpha_1 \\ \alpha_2 \\ \alpha_3 \\ \alpha_4 \end{bmatrix} = \begin{bmatrix} a_{11}^2\alpha_1 + a_{21}^2\alpha_2 + 2a_{11}a_{31}\alpha_3 + 2a_{21}a_{31}\alpha_4 \\ a_{12}^2\alpha_1 + a_{22}^2\alpha_2 + 2a_{12}a_{32}\alpha_3 + 2a_{22}a_{32}\alpha_4 \\ (a_{11}a_{22} + a_{21}a_{12})(a_{11}\alpha_3 + a_{21}\alpha_4) \\ (a_{11}a_{22} + a_{21}a_{12})(a_{12}\alpha_3 + a_{22}\alpha_4) \end{bmatrix},$$

$$\phi \begin{bmatrix} \alpha'_1 \\ \alpha'_2 \\ \alpha'_3 \\ \alpha'_4 \end{bmatrix} = \begin{bmatrix} a_{11}^2\alpha'_1 + a_{21}^2\alpha'_2 + 2a_{11}a_{31}\alpha'_3 + 2a_{21}a_{31}\alpha'_4 \\ a_{12}^2\alpha'_1 + a_{22}^2\alpha'_2 + 2a_{12}a_{32}\alpha'_3 + 2a_{22}a_{32}\alpha'_4 \\ (a_{11}a_{22} + a_{21}a_{12})(a_{11}\alpha'_3 + a_{21}\alpha'_4) \\ (a_{11}a_{22} + a_{21}a_{12})(a_{12}\alpha'_3 + a_{22}\alpha'_4) \end{bmatrix}.$$

We need $(\alpha_3, \alpha_4) \neq (0,0)$ or $(\alpha'_3, \alpha'_4) \neq (0,0)$ in order to have that c do not lie in $\theta_1^\perp \cap \theta_2^\perp$. Without loss of generality we can assume that $(\alpha_3, \alpha_4) \neq (0,0)$ moreover we can assume that $\alpha_3 \neq 0$ and $\alpha'_3 = 0$.

For θ_1 : Case (a) If $\alpha_2 = \alpha_4 = 0$, choose $a_{21} = a_{12} = 0, a_{11} = 1, a_{22} = \frac{1}{\alpha_3}, a_{31} = \frac{-\alpha_1}{2\alpha_3}$. Then we can assume that $\theta_1 = \delta_{a,c}$. Case (b) If $\alpha_2 \neq 0, \alpha_4 = 0$, choose $a_{21} = a_{12} = 0, a_{11} = \sqrt{\frac{\alpha_2}{\alpha_3}}, a_{22} = \frac{1}{\sqrt{\alpha_2}}, a_{31} = \frac{-\alpha_1}{2\alpha_3}\sqrt{\frac{\alpha_2}{\alpha_3}}$. Then we can assume that $\theta_1 = \delta_{b,b} + \delta_{a,c}$. Case (c) If $\alpha_4 \neq 0$, choose $a_{21} = a_{12} = 0, a_{11} = \sqrt{\frac{\alpha_4}{\alpha_3^2}}, a_{22} = \sqrt{\frac{\alpha_3}{\alpha_4^2}}, a_{31} = \frac{-\alpha_1}{2\alpha_3}\sqrt{\frac{\alpha_4}{\alpha_3^2}}, a_{31} = \frac{-\alpha_2}{2\alpha_4}\sqrt{\frac{\alpha_3}{\alpha_4^2}}$. Then we can assume that $\theta_1 = \delta_{a,c} + \delta_{b,c}$.

For θ_2 : Case (1) Suppose that $\theta_1 = \delta_{a,c}$: Now to fix θ_1 (up to a scalar), we require $a_{21} = a_{12} = a_{31} = 0$. It follows that $\alpha'_1 \longrightarrow a_{11}^2\alpha'_1, \alpha'_2 \longrightarrow a_{22}^2\alpha'_2 + 2a_{22}a_{32}\alpha'_4, \alpha'_3 \longrightarrow 0, \alpha'_4 \longrightarrow a_{11}a_{22}^2\alpha'_4$. Case (1.1) If $\alpha'_4 \neq 0$: By taking $a_{32} = \frac{-a_{22}\alpha'_2}{2\alpha'_4}$ we get that $\alpha'_1 \longrightarrow a_{11}^2\alpha'_1, \alpha'_2 \longrightarrow 0, \alpha'_3 \longrightarrow 0, \alpha'_4 \longrightarrow a_{11}a_{22}^2\alpha'_4$. Case (1.1.1) If $\alpha'_1 = 0$: Choose $a_{11} = \frac{1}{\alpha'_4}, a_{22} = 1, a_{32} = \frac{-\alpha'_2}{2\alpha'_4}$, so we can assume that $\theta_2 = \delta_{b,c}$ then we get a subspace $V_1 = \delta_{a,c} \wedge \delta_{b,c}$. Case (1.1.2) If $\alpha'_1 \neq 0$: Choose $a_{11} = \frac{1}{\sqrt{\alpha'_1}}, a_{22} = \sqrt{\frac{\alpha'_1}{\alpha'_2}}, a_{32} = \frac{-\sqrt{\alpha'_1\alpha'_2}}{2\alpha'_4}$, so we can assume that $\theta_2 = \delta_{a,a} + \delta_{b,c}$ then we get a subspace $V_2 = \delta_{a,c} \wedge (\delta_{a,a} + \delta_{b,c})$. Case (1.2) If $\alpha'_4 = 0$: Then we have $\alpha'_1 \longrightarrow a_{11}^2\alpha'_1, \alpha'_2 \longrightarrow a_{22}^2\alpha'_2, \alpha'_3 \longrightarrow 0, \alpha'_4 \longrightarrow 0$. Case (1.2.1) If $\alpha'_1 \neq 0, \alpha'_2 = 0$: Choose $a_{11} = \frac{1}{\sqrt{\alpha'_1}}$, so we can assume that $\theta_2 = \delta_{a,a}$ then we get a subspace

$V_3 = \delta_{a,c} \wedge \delta_{a,a}$. Case (1.2.2) If $\alpha_1' = 0, \alpha_2' \neq 0$: Choose $a_{22} = \frac{1}{\sqrt{\alpha_2'}}$, so we can assume that $\theta_2 = \delta_{b,b}$ then we get a subspace $V_4 = \delta_{a,c} \wedge \delta_{b,b}$. Case (1.2.3) If $\alpha_1' \neq 0, \alpha_2' \neq 0$: Choose $a_{11} = \frac{1}{\sqrt{\alpha_1'}}, a_{22} = \frac{1}{\sqrt{\alpha_2'}}$, so we can assume that $\theta_2 = \delta_{a,a} + \delta_{b,b}$ then we get a subspace $V_5 = \delta_{a,c} \wedge (\delta_{a,a} + \delta_{b,b})$.

Case (2) Suppose that $\theta_1 = \delta_{b,b} + \delta_{a,c}$: Now to fix θ_1 (up to a scalar), we requirea $_{21} = a_{12} = a_{31} = 0, a_{22} = a_{11}^2$. It follows that $\alpha_1' \longrightarrow a_{11}^2\alpha_1', \alpha_2' \longrightarrow a_{11}^4\alpha_2' + 2a_{11}^2 a_{32}\alpha_4', \alpha_3' \longrightarrow 0, \alpha_4' \longrightarrow a_{11}^5\alpha_4'$. Case (2.1) If $\alpha_4' \neq 0$: We can solve for a_{32} to have $\alpha_2' \longrightarrow 0$, so we can assume that $\alpha_2' = 0$. Choose $a_{32} = 0$ to keep $\alpha_2' \longrightarrow 0$, then $\alpha_1' \longrightarrow a_{11}^2\alpha_1', \alpha_2' \longrightarrow 0, \alpha_3' \longrightarrow 0, \alpha_4' \longrightarrow a_{11}^5\alpha_4'$. Case (2.1.1) If $\alpha_1' \neq 0$: Choose $a_{11} = \sqrt[3]{\frac{\alpha_1'}{\alpha_4'}}$ then $\alpha_1' \longrightarrow \alpha_1' \sqrt[3]{\frac{\alpha_1'^2}{\alpha_4'^2}}, \alpha_2' \longrightarrow 0, \alpha_3' \longrightarrow 0, \alpha_4' \longrightarrow \alpha_1' \sqrt[3]{\frac{\alpha_1'^2}{\alpha_4'^2}}$, so after dividing we can assume that $\theta_2 = \delta_{a,a} + \delta_{b,c}$. Then we get a subspace $V_6 = (\delta_{b,b} + \delta_{a,c}) \wedge (\delta_{a,a} + \delta_{b,c})$. Case (2.1.2) If $\alpha_1' = 0$: After dividing we can assume that $\theta_2 = \delta_{b,c}$, then we get a subspace $V_7 = (\delta_{b,b} + \delta_{a,c}) \wedge \delta_{b,c}$. Case (2.2) If $\alpha_4' = 0$: We have $\alpha_1' \longrightarrow a_{11}^2\alpha_1', \alpha_2' \longrightarrow a_{11}^4\alpha_2', \alpha_3' \longrightarrow 0, \alpha_4' \longrightarrow 0$. Case (2.2.1) If $\alpha_1' \neq 0, \alpha_2' \neq 0$: Choose $a_{11} = \sqrt{\frac{\alpha_1'}{\alpha_2'}}$ then $\alpha_1' \longrightarrow \frac{\alpha_1'^2}{\alpha_2'}, \alpha_2' \longrightarrow 0, \alpha_3' \longrightarrow 0, \alpha_4' \longrightarrow \frac{\alpha_1'^2}{\alpha_2'}$, so after dividing we can assume that $\theta_2 = \delta_{a,a} + \delta_{b,b}$. Then we get a subspace $V_8 = (\delta_{b,b} + \delta_{a,c}) \wedge (\delta_{a,a} + \delta_{b,b})$. Case (2.2.2) If $\alpha_1' \neq 0, \alpha_2' = 0$: After dividing we can assume that $\theta_2 = \delta_{a,a}$, then we get a subspace $V_9 = (\delta_{b,b} + \delta_{a,c}) \wedge \delta_{a,a}$. Case (2.2.3) If $\alpha_1' = 0, \alpha_2' \neq 0$: After dividing we can assume that $\theta_2 = \delta_{b,b}$ then after subtracting it from θ_1 we get $\theta_1 = \delta_{a,c}$. We are back in case (1.2.2).

Case (3) Suppose that $\theta_1 = \delta_{a,c} + \delta_{b,c}$: Now to fix θ_1 (up to a scalar), we requirea $a_{31} = a_{32} = 0$ and $a_{11} = a_{22}, a_{12} = a_{21} = 0$ or $a_{12} = a_{21}, a_{11} = a_{22} = 0$. Case (3.1) If $\alpha_1' \neq 0, \alpha_4' \neq 0$: Choosee $a_{11} = \frac{\alpha_1'}{\alpha_4'}, a_{12} = a_{21} = 0$ then after dividing by $\frac{\alpha_1'^3}{\alpha_4'^2}$ we can assume $\alpha_1' = \alpha_4' = 1$. So we can assume that $\theta_2 = \delta_{a,a} + \alpha\delta_{b,b} + \delta_{b,c}$ then we get a subspaces $V_{10}^{\alpha} = (\delta_{a,c} + \delta_{b,c}) \wedge (\delta_{a,a} + \alpha\delta_{b,b} + \delta_{b,c})$. Case (3.2) $\alpha_2' \neq 0, \alpha_4' \neq 0$: Choosee $a_{11} = a_{22} = 0$ then we are back in case (3.1). Case (3.3) $\alpha_1' \neq 0, \alpha_4' = 0$: After dividing we can assume that $\alpha_1' = 1$, then $\theta_2 = \delta_{a,a} + \alpha\delta_{b,b}$. So we get a subspaces $V_{11}^{\alpha} = (\delta_{a,c} + \delta_{b,c}) \wedge (\delta_{a,a} + \alpha\delta_{b,b})$. Case (3.4) $\alpha_2' \neq 0, \alpha_4' = 0$: Choosee $a_{11} = a_{22} = 0$ then we are back in case (3.3). Case (3.4) $\alpha_1' = 0, \alpha_4' \neq 0$: If $\alpha_2' \neq 0$ choosee $a_{11} = a_{22} = 0$ then we are back in case (3.1), If $\alpha_2' = 0$ then after dividing we can assume that $\theta_2 = \delta_{b,c}$. Subtracting θ_2 from θ_1 we get $\theta_1 = \delta_{a,c}$. Hence we are back in case (1.1.1).

Now let $J_{V_2} \xrightarrow{\varphi_1} J_{V_7} \xrightarrow{\varphi_2} J_{V_{10}^{\alpha=0}}$ such that $\varphi_1 : a \longrightarrow b, b \longrightarrow a, c \longrightarrow c, d \longrightarrow d, e \longrightarrow e$ and $\varphi_2 : a \longrightarrow b + \frac{1}{2}c, b \longrightarrow -a, c \longrightarrow -c - \frac{1}{2}d, d \longrightarrow -e, e \longrightarrow d + e$ then $J_{V_2} \cong J_{V_7} \cong J_{V_{10}^{\alpha=0}}$. Next let $J_{V_5} \xrightarrow{\varphi_3} J_{V_8}$ & $J_{V_6} \xrightarrow{\varphi_4} J_{V_{10}^{\alpha\neq 0}}$ such that $\varphi_3 : a \longrightarrow a + \frac{1}{2}c, b \longrightarrow b, c \longrightarrow c, d \longrightarrow d, e \longrightarrow -d + e$ and $\varphi_4 : a \longrightarrow \frac{1}{\alpha^{\frac{2}{3}}}a + \frac{1}{2\alpha}c, b \longrightarrow -\frac{1}{\alpha^{\frac{1}{3}}}b - \frac{1}{2}c, c \longrightarrow -\frac{1}{\alpha}c - \frac{1}{2\alpha^{\frac{2}{3}}}d - \frac{1}{2\alpha^{\frac{4}{3}}}e, d \longrightarrow -\frac{1}{\alpha^{\frac{5}{3}}}d, e \longrightarrow \frac{1}{\alpha^{\frac{5}{3}}}d + \frac{1}{\alpha^{\frac{4}{3}}}e$ then $J_{V_5} \cong J_{V_8}$ and $J_{V_6} \cong J_{V_{10}^{\alpha\neq 0}}$.

A Gröbner basis computation shows that $V_1, V_2, .., V_6, V_9$ are not in the same $Aut(J_{3,3})$-orbit and V_{11}^{α} is not be $Aut(J_{3,3})$-conjugate to $V_1, V_2, .., V_6, V_9$. Finally $J_{V_{11}^{\alpha}}$ and $J_{V_{11}^{\beta}}$ are isomorphic if and only if $\beta(\alpha^2+1) = \alpha(\beta^2+1)$. i.e, V_{11}^{α} and V_{11}^{β} lie in the same $Aut(J_{3,3})$-orbit if and only if $\beta = \alpha$ or $\beta = \alpha^{-1}$. Hence we get algebras :

- $J_{5,37} : a \circ b = c, a \circ c = d, b \circ c = e.$
- $J_{5,38} : a \circ b = c, a \circ c = d, a^2 = e, b \circ c = e.$
- $J_{5,39} : a \circ b = c, a \circ c = d, a^2 = e.$
- $J_{5,40} : a \circ b = c, a \circ c = d, b^2 = e.$
- $J_{5,41} : a \circ b = c, a \circ c = d, a^2 = e, b^2 = e.$
- $J_{5,42} : a \circ b = c, b^2 = d, a \circ c = d, a^2 = e, b \circ c = e.$
- $J_{5,43} : a \circ b = c, b^2 = d, a \circ c = d, a^2 = e.$
- $J_{5,44}^{\alpha} : a \circ b = c, a \circ c = d, b \circ c = d, a^2 = e, b^2 = \alpha e.$

Remark 4.5. $J_{5,44}^{\alpha}$ is isomorphic to $J_{5,44}^{\beta}$ if and only if $\beta = \alpha$ or $\beta = \alpha^{-1}$. In the latter case, $\varphi(a) = \alpha b', \varphi(b) = \alpha a', \varphi(c) = \alpha^2 c', \varphi(d) = \alpha^3 d', \varphi(e) = \alpha e'$, defines an isomorphism $\varphi : J_{5,44}^{\alpha} \longrightarrow J_{5,44}^{\alpha^{-1}}$.

5 Main results

In this section we summarize our main results in the following theorem.

Theorem 5.1 (**Classification of five dimensional nilpotent Jordan algebras**). *There are infinite number of five dimensional nilpotent Jordan algebras over an algebraically closed fields of characteristic $\neq 2$, and any nilpotent non-associative Jordan algebra of dimension five over an algebraically closed fields of characteristic $\neq 2$ is isomorphic to one of the following algebras :*

- $J_{5,1} : a^2 = b, b \circ c = d.$
- $J_{5,2} : a^2 = b, b \circ c = d, b \circ d = e.$
- $J_{5,3} : a^2 = b, b \circ c = d, b \circ d = e, c^2 = e.$
- $J_{5,4} : a \circ b = c, a \circ c = d.$
- $J_{5,5} : a \circ b = c, a \circ c = d, b^2 = d.$

- $J_{5,6} : a \circ b = c, a \circ c = d, b \circ c = d.$
- $J_{5,7} : a^2 = b, d^2 = e, b \circ c = e.$
- $J_{5,8} : a^2 = b, a \circ d = e, b \circ c = e.$
- $J_{5,9} : a \circ b = c, c \circ d = e.$
- $J_{5,10} : a \circ b = c, c \circ d = e, a^2 = e.$
- $J_{5,11} : a \circ b = c, c \circ d = e, a^2 = e, b^2 = e.$
- $J_{5,12} : a \circ b = c, a \circ c = e, d^2 = e, b \circ c = e.$
- $J_{5,13} : a \circ b = c, a \circ c = e, d^2 = e.$
- $J_{5,14} : a \circ b = c, a \circ c = e, d^2 = e, b^2 = e.$
- $J_{5,15} : a \circ b = c, a \circ c = e, a \circ d = e, b \circ c = e.$
- $J_{5,16} : a \circ b = c, a \circ c = e, b \circ d = e.$
- $J^{\alpha}_{5,17} : a^2 = b, a \circ b = c, a \circ c = e, b^2 = e, b \circ d = e, d^2 = \alpha e.$
- $J_{5,18} : a \circ b = d, c^2 = d, a \circ d = e.$
- $J_{5,19} : a \circ b = d, c^2 = d, a \circ d = e, b \circ c = e.$
- $J_{5,20} : a \circ b = d, c^2 = d, a \circ d = e, b^2 = e.$
- $J_{5,21} : a \circ b = d, c^2 = d, c \circ d = e.$
- $J_{5,22} : a \circ b = d, c^2 = d, c \circ d = e, a^2 = e.$
- $J_{5,23} : a \circ b = d, c^2 = d, c \circ d = e, a^2 = e, b^2 = e.$
- $J_{5,24} : a^2 = b, a \circ b = d, c^2 = d, b^2 = e, a \circ d = e.$
- $J_{5,25} : a^2 = c, a \circ b = d, b \circ c = e, b \circ d = e.$
- $J^{\alpha}_{5,26} : a^2 = c, a \circ b = d, a \circ c = e, b \circ d = e, b \circ c = \alpha e.$
- $J^{\alpha \neq 0,1}_{5,27} : a^2 = c, a \circ b = d, b \circ c = e, a \circ d = \alpha e.$
- $J_{5,28} : a^2 = c, a \circ b = d, a \circ c = e, b \circ c = -2e, a \circ d = e.$
- $J^{\alpha}_{5,29} : a^2 = c, b^2 = d, a \circ c = e, a \circ d = e, b \circ d = \alpha e.$

- $J^{\alpha,\beta}_{5,30} : a^2 = c, b^2 = d, b \circ c = e, a \circ d = e, b \circ d = \alpha e, a \circ c = \beta e.$
- $J_{5,31} : a^2 = b, b \circ c = d, a \circ b = e.$
- $J_{5,32} : a^2 = b, b \circ c = d, a \circ b = e, a \circ c = e.$
- $J_{5,33} : a^2 = b, b \circ c = d, a \circ b = e, c^2 = e.$
- $J_{5,34} : a^2 = b, b \circ c = d, c^2 = e.$
- $J_{5,35} : a^2 = b, b \circ c = d, a \circ c = e.$
- $J_{5,36} : a^2 = b, b \circ c = d, a \circ c = e, c^2 = e.$
- $J_{5,37} : a \circ b = c, a \circ c = d, b \circ c = e.$
- $J_{5,38} : a \circ b = c, a \circ c = d, a^2 = e, b \circ c = e.$
- $J_{5,39} : a \circ b = c, a \circ c = d, a^2 = e.$
- $J_{5,40} : a \circ b = c, a \circ c = d, b^2 = e.$
- $J_{5,41} : a \circ b = c, a \circ c = d, a^2 = e, b^2 = e.$
- $J_{5,42} : a \circ b = c, b^2 = d, a \circ c = d, a^2 = e, b \circ c = e.$
- $J_{5,43} : a \circ b = c, b^2 = d, a \circ c = d, a^2 = e.$
- $J^{\alpha}_{5,44} : a \circ b = c, a \circ c = d, b \circ c = d, a^2 = e, b^2 = \alpha e.$

where $\alpha, \beta \in \Bbbk$.
Among these algebras there are precisely the following isomorphisms :

- $J^{\alpha}_{5,26} \approxeq J^{\beta}_{5,26}$ *if and only if* $\alpha^2 - \beta^2 = 0$.
- $J^{\alpha}_{5,29} \approxeq J^{\beta}_{5,29}$ *if and only if* $\alpha^2 - \beta^2 = 0$.
- $J^{\alpha,\beta}_{5,30} \approxeq J^{\alpha',\beta'}_{5,30}$ *if and only if* $(\alpha, \beta) = (\alpha', \beta')$ *or* $(\alpha, \beta) = (\beta', \alpha')$.
- $J^{\alpha}_{5,44} \approxeq J^{\beta}_{5,44}$ *if and only if* $\beta = \alpha$ *or* $\beta = \alpha^{-1}$.

References

[1] Ahmed Hegazi and Hani Abdelwahab. Construction of nilpotent Jordan algebras over any arbitrary fields. arXiv:1301.6349 [math.RA], 2013.

[2] Guerino Mazzola. Generic finite schemes and Hochschild cocycles. *Comment. Math. Helv.*, 55(2):267–293, 1980.

[3] Bjorn Poonen. Isomorphism types of commutative algebras of finite rank over an algebraically closed field. In *Computational arithmetic geometry*, volume 463 of *Contemp. Math.*, pages 111–120. Amer. Math. Soc., Providence, RI, 2008.

[4] W. Bosma, J. Cannon, and C. Playoust. The Magma algebra system I: The user language. *J. Symbolic Comput.*, 24(3-4):235–265, 1997.

[5] A. A. Albert. A structure theory for Jordan algebras. *Ann.Math.* 48 (1946),546-567.

[6] Schafer, Richard D. .An introduction to nonassociative algebras. *Pure and Applied Mathematics*, Vol. 22, *Academic Press*, New York, 1966.

[7] W. A. de Graaf. Classification of 6-dimensional nilpotent Lie algebras over fields of characteristic not 2. *J. Algebra*, 309:640–653, 2007.

[8] Tor Skjelbred and Terje Sund. Sur la classification des algèbres de Lie nilpotentes. *C. R. Acad. Sci. Paris Sér. A-B*, 286(5), 1978.

[9] D. Cox, J. Little, and D. O'Shea. *Ideals, Varieties and Algorithms: An Introduction to Computational Algebraic Geometry and Commutative Algebra*. Springer Verlag, New York, Heidelberg, Berlin, 1992.

[10] R.E. Beck, B. Kolman. Construction of nilpotent Lie algebras over arbitrary fields. In Paul S. Wang, editor, *Proceedings of the 1981 ACM Symposium on Symbolic and Algebraic Computation*, pages 169–174. ACM New York, 1981.

[11] M.-P. Gong. *Classification of Nilpotent Lie Algebras of Dimension 7*. PhD thesis, University of Waterloo, Waterloo, Canada, 1998.